An Introduction to Asset Pricing Theory

Junhui Qian

Preface

This book is derived from my lecture notes for an advanced course on asset pricing theory. The purpose is to provide a concise and accessible introduction to no-arbitrage asset pricing for students who have taken undergraduate courses in calculus, probability and statistics, and linear algebra.

The readers may include advanced undergraduate and graduate students in finance, economics, and applied mathematics, as well as researchers and practitioners in the field of quantitative finance.

The book is concise, making it ideal for students and practitioners who are looking for a quick introduction to the topic. I often sacrifice rigor to make the book easier to read and understand. However, I have taken care to include all key steps in the proofs and derivations to ensure that readers can follow along.

I hope that this book will serve as a helpful guide to the fundamentals of noarbitrage asset pricing and will help readers appreciate the beauty of mathematical finance.

Shanghai, China, April 2024 Junhui Qian junhuiq@gmail.com

Contents

Preface i								
	1							
	2							
	8							
	13							
	13							
	14							
	14							
	15							
	16							
	18							
	18							
	18							
	21							
	24							
	24							
	26							
	29							
	31							
•	 							

3 No-Arbitrage Pricing in Continuous Time

	3.1	Basic Setup	41
	3.2	The Black-Scholes Model	44
	3.3	The Feynman-Kac Solution	46
	3.4	Risk-Neutral Pricing	49
	3.5	State Prices	59
	3.6	Treatment of Dividends	61
4	Ter	m Structure Modeling	65
	4.1	Basics	65
	4.2	The Single-Factor Heath-Jarrow-Morton Model $\ . \ . \ . \ . \ .$.	68
	4.3	Short-Rate Models	71
	4.4	Multi-factor Models	77
	4.5	Pricing Interest Rate Products	80
	4.6	Forward Measure	83
A	App	pendix to Chapter 1	89
	A.1	Classical Derivation of CAPM	89
		A.1.1 Efficiency Frontier without Riskfree Asset	89
		A.1.2 CAPM	90

Chapter 1

Introduction

The theory of asset pricing is concerned with explaining and determining prices of financial assets in a uncertain world.

The asset prices we discuss would include prices of stocks, bonds, exchange rates, and derivatives of all these underlying financial assets. Asset pricing is crucial for the allocation of financial resources. Mispricing of financial assets would lead to inefficiency in investment and consumption in the real economy.

The "uncertainty" in this book is, rather simplistically, described by probability distributions. A more sophisticated treatment would differentiate uncertainty from risk as in Knight (1921). Here we treat uncertainty and risk as the same thing: future variation that can be characterized by some distribution without ambiguity. In this book, uncertainty is assumed in both how an asset would pay in the future and how agents would discount the payoff.

In this book we also take the simplistic view that the uncertainty is given and that it is not influenced by the evolution of prices. It is generally believed in the investment community, however, that prices may affect future payoffs. For example, a surge in stock price would lower financing cost for the company and boost future earnings. We do not go into this direction.

In this first chapter, we study no-arbitrage pricing in a two-period world. Key concepts such as state prices, risk-neutral probability, and stochastic discount factor, are introduced.

Classical asset pricing models, such as CAPM and APT (Arbitrage Pricing Theory), are discussed as special cases of modern asset pricing theory using stochastic discount factor. A classical derivation of CAPM is offered in the Appendix.

1.1 No-Arbitrage Pricing in Finite States

We first introduce some basic concepts of asset pricing in a two-period finite-state world.

Commodity A commodity is a "good" at a particular time and a particular place when a particular "state" happens. For a commodity to be interesting to economists, it must cost something. Besides physical characteristics, the key characteristic of a commodity is its availability in space and time, and conditionality. A cup of water in the desert and another one in Shanghai, although physically the same, are different commodities. And an umbrella when it rains is also different from that when it does not. It is in the sense of conditionality that we call a commodity a "contingent claim". For example, forwards and futures on oil, ores, gold, and other metals can be understood as commodities.

Security Security is financial commodity. Stocks, bonds, and their derivatives are all securities. The payoffs of physical goods are physical goods. The payoffs of securities are mostly money, sometimes other securities. A financial market is a market where securities are exchanged.

Consider a two-period world. We live at time t, the next period is t + 1. The essential characteristics of a security in this world include the price p_t and the future payoff x_{t+1} . At time t, p_t is observed and x_{t+1} are a random variable.

For examples, we understand

- stock: $x_{t+1} = p_{t+1} + d_{t+1}$, where d_{t+1} denotes dividend payment.
- bond (zero-coupon, riskless): $x_{t+1} = 1$.
- forwards/futures (on a stock with strike price K, long position):

$$x_{t+1} = p_{t+1} - K$$

• option (European, on a stock with strike price K)

long call:
$$x_{t+1} = \max\{0, p_{t+1} - K\}$$

long put: $x_{t+1} = \max\{0, K - p_{t+1}\}.$

For now, we assume x_{t+1} is a discrete finite-state random variable, taking value in \mathbb{R}^S . In other words, there are S possible values for x_{t+1} , $(x^1, x^2, \dots, x^S)'$, corresponding to states $s = 1, 2, \dots, S$ and probability $\pi = (\pi^1, \pi^2, \dots, \pi^S)'$. We assume $\pi^s > 0$ for all s.

Portfolio Suppose there are J securities in the market, with prices described by a vector $p = (p_1, p_2, \ldots, p_J)'$, where p_j is the price of security j. Individuals may build portfolios of these securities. Mathematically, a portfolio is characterized by a J-dimensional vector $h \in \mathbb{R}^J$. The total price of a portfolio h is thus p'h.

We use a matrix to describe payoffs of all securities at time t + 1:

$$X = \begin{pmatrix} x_1' \\ x_2' \\ \vdots \\ x_J' \end{pmatrix} = \begin{pmatrix} x_1^1 & x_1^2 & \cdots & x_1^S \\ x_2^1 & x_2^2 & \cdots & x_2^S \\ \vdots & & & \\ x_J^1 & x_J^2 & \cdots & x_J^S \end{pmatrix}$$

Note that rows correspond to securities and columns correspond to states. The payoff of a portfolio h at time t + 1 is thus, X'h.

Complete market If rank(X) = S, the financial market is complete, meaning that every payoff vector in \mathbb{R}^S can be realized by trading these J securities. More formally, for all $x \in \mathbb{R}^S$, there exists $h \in \mathbb{R}^J$ such that x = X'h.

Asset span For any financial market, the asset span is the space spanned by columns of X':

$$\mathcal{M} = \{ X'h, h \in \mathbb{R}^J \} = \operatorname{span}(X').$$

If $\mathcal{M} = \mathbb{R}^S$, the market is complete.

The Law of One Price The law of one price (LOP) states that portfolios with the same payoff must have the same price:

$$X'h = X'\tilde{h} \quad \Rightarrow \quad p'h = p'\tilde{h},$$

where $p \in \mathbb{R}^J$ is the price vector.

Theorem A necessary and sufficient condition for LOP is: zero payoff has zero price.

Proof : (1) If LOP holds, $X'(h-\tilde{h}) = 0 \implies p'(h-\tilde{h}) = 0$. (2) If LOP does not hold, then $X'h = X'\tilde{h}$ but $p'h \neq p'\tilde{h}$, this is, there exists a h^* such that $X'h^* = 0$ but $p'h^* \neq 0$.

Theorem For any $z \in \mathcal{M}$, there is a linear pricing functional¹ q(z) if and only if LOP holds.

Proof (1) Linear functional \Rightarrow LOP holds. If LOP does not hold, then $q(0) = \epsilon \neq 0$, then $q(z+0) = q(z) + \epsilon$, contradicting the definition of linear functional. (2) LOP holds \Rightarrow linearity. For any $z, \tilde{z} \in \mathcal{M}$, we can find h and \tilde{h} such that z = X'h, $\tilde{z} = X'\tilde{h}$. So $\alpha z + \beta \tilde{z} = \alpha X'h + \beta X'\tilde{h}$. So

$$q(\alpha z + \beta \tilde{z}) = \alpha p'h + \beta p'\tilde{h} = \alpha q(z) + \beta q(\tilde{z}).$$

No Arbitrage For a vector $x = (x_1, \ldots, x_n)'$, we define

 $x \ge 0$ if $x_i \ge 0$ for all i,

x > 0 if $x_i \ge 0$ for all i and $x_i > 0$ for some i,

Arbitrage An arbitrage is a portfolio h that satisfies $X'h \ge 0$ and p'h < 0. An arbitrage portfolio generates nonnegative payoff but has a negative price. If there are arbitrage opportunities, the market is obviously not stable or efficient. The no-arbitrage assumption is thus a weak form of equilibrium or efficiency.

Theorem The payoff functional is linear and positive if and only if there is no arbitrage.

Proof (1) linear & positivity \Rightarrow no arbitrage: For any h that satisfies $X'h \ge 0$, $p'h = q(X'h) \ge 0$. (2) no arbitrage \Rightarrow linear & positivity: no arbitrage \Rightarrow LOP \Rightarrow linearity, and positivity follows from $z = X'h \ge 0 \Rightarrow q(z) = p'h \ge 0$, for any h.

Note that a functional is positive if it assigns nonnegative value to every positive element of its domain. It is strictly positive if it assigns strictly positive value to positive elements.

An Example Let's have some flavor of no-arbitrage pricing. Consider a financial market with a money account, a stock, and an European call option on the stock with strike price 98. Suppose there are two future states. If state 1 realizes, the stock price declines to 84 from the current price 100. If state 2 happens, the stock price rises to 112. Suppose the interest rate on the money account is 5%, we want to obtain a no-arbitrage price c_0 for the call option. The following table lists the payoff structure of our simple financial market.

¹A functional is a mapping from a vector space, which may be infinite-dimensional, to real scalar numbers.

	Time 0	Time 1	
		State 1	State 2
Money	1	1.05	1.05
Stock	100	84	112
Call	c_0	0	(112-98)=14

The idea of no-arbitrage pricing is to form a portfolio of Money and Stock, $h = (\alpha, \beta)'$, that replicates the payoff of the call option, and to deduce the option price from the current price of the replication portfolio. If there is no arbitrage opportunities in the market, then the price of option must be the same as that of the replication portfolio.

We solve the following set of equations with unknown α and β ,

$$1.05\alpha + 84\beta = 0$$

$$1.05\alpha + 112\beta = 14,$$

and obtain,

$$\alpha = -40 \quad \beta = 1/2.$$

This means that we borrow 40 from bank and buy a "half" stock. The portfolio h = (-40, 1/2)' exactly replicates the payoff of the call option. At time 0, this portfolio has a value of

$$100 \cdot \frac{1}{2} - 40 = 10.$$

And this should be the price of the call option.

If the option price is 5, then we can form the following portfolio

$$(\alpha, \beta, c) = (45, -1/2, 1)$$

The current price of the portfolio is zero, but the payoff will be 5.25 next period whichever state realizes. This is an arbitrage. If the price is 15, readers may verify that the following portfolio achieves an arbitrage

$$(\alpha, \beta, c) = (-35, 1/2, -1).$$

Obviously, short selling must be allowed to make the above analysis valid. Borrowing from banks can be considered as shorting the money.

State prices Consider a special type of securities with following payoff,

$$e^s = (\underbrace{0, 0, \cdots, 0}_{S}, 1, 0, \cdots, 0_{S})'.$$

When and only when state s happens, the security gives one unit of payoff. Economists call them Arrow-Debreu securities. In a complete market with S states, all of the S Arrow-Debreu securities should be available.

Using Arrow-Debreu securities, we can represent any asset payoff x by a portfolio:

$$x = (x^1, \cdots, x^S)' = \sum_{s=1}^{S} x^s e^s.$$

Let $\varphi^s = q(e^s)$, and

$$\varphi = (\varphi^1, \varphi^2, \cdots, \varphi^S)'$$

 φ is called the vector of state prices. The no-arbitrage price of x would be

$$q(x) = \sum_{s=1}^{S} x^s q(e^s) = \sum_{s=1}^{S} x^s \varphi^s = x' \varphi.$$

Theorem There is no arbitrage if and only if there is a state price vector. To prove this theorem, we need the Stiemke's Lemma, which is a theorem of alternatives.

The Stiemke's Lemma For an m-by-n matrix A, one and only one of the following statements is true:

- (a) There exists an $x \in \mathbb{R}^n$ and x > 0 such that Ax = 0.
- (b) There exists a $y \in \mathbb{R}^m$ with y'A > 0.

Recall that an arbitrage-portfolio is one that satisfies $X'h \ge 0$ and p'h < 0. This can be stated mathematically,

$$h'(-p,X) > 0.$$

According to the Stiemke's Lemma, there is no such h if and only if there exists a vector $\varphi \in \mathbb{R}^S$ and $\varphi > 0$ such that

$$(-p,X)\left(\begin{array}{c}1\\\varphi\end{array}\right)=0.$$

Put differently, we have

$$p = X\varphi.$$

The vector φ is the desired state price vector. To see this, let X be an S-by-S identity matrix, which characterizes a market for S Arrow-Debreu securities. We then have $p = \varphi$, which means that $q(e^s) = \varphi^s$ for all s.

Example (continued) In the previous example (the market with a money account and a stock), the payoff matrix and the price vector are as follows,

$$X = \begin{bmatrix} 1.05 & 1.05 \\ 84 & 112 \end{bmatrix}, \quad p = \begin{bmatrix} 1 \\ 100 \end{bmatrix}.$$

The state price φ can be determined by solving

$$X\varphi = p,$$

which yields

$$\varphi = \left[\begin{array}{c} 0.2381\\ 0.7143 \end{array} \right].$$

Using the state price vector, we can price the European call option:

$$c_0 = 0.2381 \times 0 + 0.7143 \times 14 \approx 10.$$

Risk-neutral probability Let

$$\varphi^0 = \sum_{s=1}^S \varphi^s = \sum_{s=1}^S q(e^s) = q(\iota),$$

where ι is the vector of 1's. Note that φ^0 is the price of risk-free zero-coupon bond with the following yield,

$$R_f = \frac{1}{\varphi^0}.$$

The price of x can be written as

$$p = q(x) = \sum_{s=1}^{S} \varphi^s x^s = \varphi^0 \sum_{s=1}^{S} \frac{\varphi^s}{\varphi_0} x^s.$$

Define

$$\tilde{\pi}^s = \frac{\varphi^s}{\varphi^0}.$$

We have

$$\tilde{\pi}^s > 0 \ \forall s \quad \text{and} \quad \sum_{s=1}^S \tilde{\pi}^s = 1.$$

We call $(\tilde{\pi}^s)$ risk-neutral probabilities. Using $(\tilde{\pi}^s)$, we can represent asset price as

$$p = \varphi^0 \sum_{s=1}^S \tilde{\pi}^s x^s = R_f^{-1} \tilde{\mathbb{E}} x,$$

where the expectation $\tilde{\mathbb{E}}$ is taken with respect to the risk-neutral probability.

Example: continued. Let the risk-neutral probability of state 1 be \tilde{p} , then the stock price should satisfy

$$100 = \frac{1}{1.05} (84\tilde{p} + 112(1 - \tilde{p})).$$

So $\tilde{p} = 1/4$. So the price of call option is

$$c_0 = \frac{1}{1.05} (0 \cdot \tilde{p} + 14 \cdot (1 - \tilde{p})) = \frac{1}{1.05} \cdot 14 \cdot \frac{3}{4} = 10.$$

Stochastic discount factor (SDF) Suppose that the objective probability of state s is π^s . We can represent the asset price as

$$p = R_f^{-1} \sum_{s=1}^S \pi^s \frac{\tilde{\pi}^s}{\pi^s} x^s.$$

Define $m^s = R_f^{-1} \frac{\tilde{\pi}^s}{\pi^s}$, $m^s > 0$. And let *m* be a random variable taking value m^s if the state *s* realizes. We have

$$p = \sum_{s=1}^{S} \pi^{s}(m^{s}x^{s})$$
$$= \mathbb{E}(mx).$$
(1.1)

m is called the *stochastic discount factor* (SDF).

1.2 Extension to Continuous States

The pricing formula in (1.1) can be extended to the world of continuous payoffs, which may be represented by continuous-state random variables. Recall that a continuous-state random variable is defined as a mapping from the sample space to the real line,

$$x \equiv x(\omega): \ \Omega \to \mathbb{R}$$

Similarly, the SDF m is a \mathbb{R} -valued random variable.

We define $\mathcal{M} = \{x \in \mathbb{R} : \mathbb{E}x^2 < \infty\}$. This set contains all "reasonable" payoffs. And we define inner product on \mathcal{M} as,

$$\langle x_1, x_2 \rangle = \mathbb{E}(x_1 x_2).$$

It is well known that \mathcal{M} is a Hilbert space with the above inner product. If there is no arbitrage, then q is a linear positive functional on \mathcal{M} . According to Riesz's

Representation Theorem, every bounded linear pricing functional q on \mathcal{M} can be represented in terms of the inner product,

$$q(x) = \langle x, m \rangle = \mathbb{E}(mx)$$

for some $m \in \mathcal{M}$. Since q must be positive to rule out arbitrage, m > 0 almost sure. The reverse is also true. Hence we may conclude that there is no arbitrage if and only if m > 0 almost sure.

We now use the pricing formula in (1.1) to understand some key concepts and models in finance.

Gross return The gross return of an asset with price p and payoff x is R = x/p. It must satisfy

$$1 = \mathbb{E}mR.$$

Risk-free rate If $x = \iota$, the vector of 1's, then it is the payoff of a risk-free bond. The price of the risk-free bond is $p = \mathbb{E}m$. And the risk-free return is given by

$$R_f = \frac{1}{\mathbb{E}m}.$$

Risk premium Let R = x/p denote the return to an asset with price p and payoff x. Using $p = \mathbb{E}(mx) = \operatorname{cov}(m, x) + (\mathbb{E}m)(\mathbb{E}x)$, we can represent the expected return as

$$ER = \frac{\mathbb{E}x}{p}$$
$$= \frac{1}{\mathbb{E}m} - \frac{\operatorname{cov}(m, R)}{\mathbb{E}m}$$
$$= R_f - R_f \operatorname{cov}(m, R)$$

Risk premium is defined as the expected return in excess of the risk-free rate, $\mathbb{E}R - R_f$. We have

 $\mathbb{E}R - R_f = -R_f \operatorname{cov}(m, R).$

Risk premium is proportional to cov(m, R).

 β -pricing For asset *i*, the gross return can be written as

$$\mathbb{E}R_i = R_f + \left(\frac{\operatorname{cov}(R_i, m)}{\operatorname{var}(m)}\right) \left(-\frac{\operatorname{var}(m)}{\mathbb{E}m}\right).$$

Define $\beta_{i,m} = \frac{\operatorname{COV}(R_i,m)}{\operatorname{Var}(m)}$ and $\lambda_m = -\frac{\operatorname{Var}(m)}{\mathbb{E}m}$. We can rewrite the above as

$$\mathbb{E}R_i = R_f + \beta_{i,m}\lambda_m$$

 $\beta_{i,m}$ measures the systematic risk contained in asset *i* and λ_m may be called "price of risk".

Factor models Suppose that *m* has a factor structure like

$$m = a + b'f,\tag{1.2}$$

where f is a vector of factors, b is the vector of factor loadings, and a a constant. This specification of SDF gives us a factor model of asset pricing. Without loss of generality, we assume $\mathbb{E}f = 0$. So we have $\mathbb{E}m = a = 1/R_f$.

Since $1 = \mathbb{E}(mR_i)$, we have

$$\mathbb{E}(R_i) = \frac{1}{\mathbb{E}m} - \frac{\operatorname{cov}(m, R_i)}{\mathbb{E}m} = \frac{1}{a} - \frac{\mathbb{E}(R_i f')b}{a}.$$

Let β_i be the regression coefficient of R_i on f, $\beta_i \equiv \mathbb{E}(ff')^{-1}\mathbb{E}(fR_i)$. So

$$\mathbb{E}(R_i) = \frac{1}{a} - \frac{\mathbb{E}(R_i f') \mathbb{E}(f f')^{-1} \mathbb{E}(f f') b}{a} = \frac{1}{a} - \beta' \frac{\mathbb{E}(f f') b}{a}.$$

Note that $\mathbb{E}(ff')b = \mathbb{E}mf$. If we define

$$\lambda \equiv -R_f \mathbb{E}(mf),$$

we have

$$\mathbb{E}R_i = R_f + \lambda'\beta_i. \tag{1.3}$$

This generalizes the β -pricing model.

The factor models includes the celebrated Capital Asset Pricing Model (CAPM) model and the Arbitrage Pricing Theory (APT) as special cases.

CAPM The Capital Asset Pricing Model (CAPM) model is most frequently stated as:

$$\mathbb{E}R_i = R_f + \beta_i (\mathbb{E}R_m - R_f), \qquad (1.4)$$

where R_m denotes the return on the "market portfolio". We usually proxy R_m by the return on a broad stock market index such as S&P 500. β_i captures systematic risk contained in *i*-th stock, which cannot be diversified away. And $(\mathbb{E}R_m - R_f)$ is the risk premium of the market portfolio, i.e., the price of systematic risk. The CAPM model tells us only systematic risk is priced. Bearing idiosyncratic risk, which can be diversified away using a portfolio, does not get rewarded. For interested readers, Appendix A.1 provides a classic derivation of the CAPM model using portfolio optimization.

In practice, β_i is estimated by running the following time-series regression for *i*-th stock,

$$R_{it} - R_{ft} = \alpha_i + \beta_i (R_{mt} - R_{ft}) + \varepsilon_{it}.$$

According to CAPM, α_i should be zero for all stocks and portfolios.

APT The Arbitrage Pricing Theory (APT) is a linear factor model. We assume that the asset payoff can be statistically characterized by a factor structure:

$$R_i = \mathbb{E}R_i + \beta'_i f + \varepsilon_i, \tag{1.5}$$

where f a K-by-1 vector of demeaned factors, $\mathbb{E}\varepsilon_i = 0$, $\mathbb{E}f\varepsilon_i = 0$, and $\mathbb{E}\varepsilon_i\varepsilon_j = 0$ for $i \neq j$. The price errors (ε_i) represent idiosyncratic risks that can be diversified away and hence not priced. If we assume $\sigma(m) < \infty$ and LOP holds, we have

$$q(R_i) = \mathbb{E}R_i q(\iota) + \beta'_i q(f).$$

Since $q(R_i) = 1$, $q(\iota) = 1/R_f$, we solve the above equation for $\mathbb{E}R_i$:

$$\mathbb{E}R_i = R_f + \beta'_i \lambda_i$$

where $\lambda = -R_f q(f)$. In empirical asset pricing, factor models (e.g., Fama & French three-factor model) are the main framework for analyzing cross-section returns.

Mean-variance frontier We have

$$1 = \mathbb{E}(mR_i) = \mathbb{E}m\mathbb{E}R_i + \operatorname{cov}(m, R_i)$$
$$= \mathbb{E}m\mathbb{E}R_i + \rho_{m, R_i}\sigma(m)\sigma(R_i)$$

So

$$\mathbb{E}R_i = \frac{1}{\mathbb{E}m} - \frac{1}{\mathbb{E}m}\rho_{m,R_i}\sigma(m)\sigma(R_i)$$
$$= R_f - \rho_{m,R_i}\frac{\sigma(m)}{\mathbb{E}m}\sigma(R_i).$$

Then, since $|\rho| \leq 1$,

$$|\mathbb{E}R_i - R_f| \le \frac{\sigma(m)}{\mathbb{E}m}\sigma(R_i).$$

Notice $\frac{\mathbb{E}R_i - R_f}{\sigma(R_i)}$ is the Sharpe ratio, and

$$\frac{\mathbb{E}R_i - R_f}{\sigma(R_i)} \le R_f \sigma(m).$$

Discounted cash flow valuation Consider a stock with price process p_t and dividend process d_t . Using the formula $p_t = \mathbb{E}_t m_{t+1} x_{t+1}$, we have

$$p_{t} = \mathbb{E}_{t}[m_{t+1}(p_{t+1} + d_{t+1})]$$

$$= \mathbb{E}_{t}[m_{t+1}(\mathbb{E}_{t+1}[m_{t+2}(p_{t+2} + d_{t+2})] + d_{t+1})]$$

$$\vdots$$

$$= \mathbb{E}_{t}\left(\prod_{j=1}^{n} m_{t+j}\right)p_{t+n} + \sum_{i=1}^{n} \mathbb{E}_{t}\left(\prod_{j=1}^{i} m_{t+j}\right)d_{t+i}$$

$$= \mathbb{E}_{t}\left(\prod_{j=1}^{\infty} m_{t+j}\right)p_{\infty} + \sum_{i=1}^{\infty} \mathbb{E}_{t}\left(\prod_{j=1}^{i} m_{t+j}\right)d_{t+i}$$

The first term is called the *rational bubble*. For it to exist as $n \to \infty$, p_{t+n} must also go to infinity. The second term on the last line is the discounted cash flow (DCF) value of the stock. If we assume that the dividend process is known at time t, and $m_t = 1/(1+r)$ with r > 0, then the second term reduces to the familiar

$$p_t = \sum_{i=1}^{\infty} \frac{d_{t+i}}{(1+r)^i}.$$

Further Reading

- Arrow, K.J. and Debreu, G., 1954, "Existence of an equilibrium for a competitive economy", Econometrica, 22, 265-290.
- Cochrane, J.H., 2005, Asset Pricing, Princeton University Press.
- Fama, E.F., French, K.R., 1992. "The Cross-Section of Expected Stock Returns". The Journal of Finance. 47 (2): 427-465.
- Hansen, L.P., Richard S.F., 1987, "The Role of Conditioning Information in Deducing Testable Restrictions Implied by Dynamic Asset Pricing Models," Econometrica, 55, 587-613.
- Knight, F.H., 1921, "Risk, Uncertainty, and Profit". Boston, MA: Hart, Schaffner & Marx; Houghton Mifflin Company.
- Ross, S., 1976, "The arbitrage theory of capital asset pricing". Journal of Economic Theory 13 (3): 341-360.

Chapter 2

Mathematical Background

2.1 Probability Setup

Random Variable

A random variable X is defined as a mapping from sample space Ω to \mathbb{R} with a probability measure \mathbb{P} defined on a σ -field of Ω , \mathcal{F} . The triple $(\Omega, \mathcal{F}, \mathbb{P})$ is called the *probability measure space*.

A σ -field of Ω is a collection of subsets of Ω containing Ω itself and the empty set ϕ , and closed under complements, countable unions. In mathematical language, the random variable X is a \mathcal{F} -measurable function from Ω to \mathbb{R} . Being \mathcal{F} -measurable is defined as

$$\{\omega \in \Omega | X(\omega) \le x\} \in \mathcal{F}, \ x \in \mathbb{R}.$$

Or, $X^{-1}(B) \in \mathcal{F}$ for every Borel set $B \in \mathcal{B}(\mathbb{R})$. $\mathcal{B}(\mathbb{R})$ is the smallest σ -field containing all open sets of \mathbb{R} . We can understand Borel sets as "nice" sets that we mortals can grasp, for example intervals like [a, b]. We are interested in knowing probabilities of X falling into these nice sets (e.g., $\mathbb{P}(X(\omega) \in [a, b])$). Being \mathcal{F} -measurable means that the inverse image of these nice sets, which are subsets of Ω , are elements of \mathcal{F} .

In intuitive terms, the σ -field \mathcal{F} is a collection of events. To see this, consider throwing dimes for three times, the sample space Ω is

 $\Omega = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}.$

One σ -field on this sample space is $\mathcal{F} = 2^{\Omega}$, the power set of Ω that contains all subsets of Ω including \emptyset and Ω . One element of \mathcal{F} is

$$\{HHH, HHT, HTH, HTT\}.$$

This is an event that may be called "Head appear in the first throw". For another example, the event $\{HTH, HTT\}$ may be called "First Head, second Tail". So X being \mathcal{F} -measurable means that $X^{-1}(B)$ is indeed an event in \mathcal{F} .

The σ -field generated by a random variable X, denoted $\sigma(X)$, is defined as

$$\sigma(X) = \{ X^{-1}(B) | B \in \mathbb{B}(\mathbb{R}) \},\$$

where $\mathbb{B}(\mathbb{R})$ denotes the Borel σ -field of \mathbb{R} . Roughly speaking, $\sigma(X)$ is the collection of events we may know through observing X.

2.2 Continuous-Time Stochastic Process

A stochastic process is a sequence of random variables ordered by time. We denote a stochastic process by $X = (X_t), t \in \mathcal{T}$, where \mathcal{T} is an index set. The index set \mathcal{T} can be a discrete set such as $\{1, 2, ...\}$, or a continuous set, say [0, 1]. If the index set is a continuous set, then we call X a continuous-time stochastic process.

More rigorously, X is a mapping from the product space of $\Omega \times \mathcal{T}$ to \mathbb{R} . So we may write $X_t = X_t(\omega) = X(\omega, t)$. $X(t, \cdot)$ is a random variable, and $X(\cdot, \omega)$ is a sample path or realization of X.

A filtration is a non-decreasing sequence of σ -fields ordered by time t, (\mathcal{F}_t) . Being non-decreasing means $\mathcal{F}_s \subset \mathcal{F}_t$ if s < t. Recall that a σ -field is a collection of events. The more inclusive a σ -field is, the more we may possibly know about the sample space Ω . So we can roughly think of σ -fields as information sets. A filtration is thus an ever-finer sequence of information sets.

The natural filtration of $X = (X_t)$ is defined by $\mathcal{F}_t = \sigma((X_s)_{s \leq t})$, that is, the σ -field generated by $((X_s)_{s \leq t})$, or intuitively speaking, the information contained in the stochastic process up to time t.

A stochastic process $X = (X_t)$ is said to be *adapted* to a filtration (\mathcal{F}_t) if, for every $t \ge 0$, X_t is a \mathcal{F}_t -measurable random variable.

2.2.1 Brownian Motion

A one-dimensional Brownian motion (BM) is defined as a continuous-time \mathbb{R} -valued process $W = (W_t)$ satisfying

(i) Continuous sample path almost sure (a.s.);

- (ii) Independent Gaussian increment: Let \mathcal{F}_s be the natural filtration of W_t . For any s < t, $W_t - W_s$ is independent of \mathcal{F}_s and $W_t - W_s | \mathcal{F}_s \sim N(0, t - s)$ for $t \ge s$;
- (iii) $W_0 = 0$ a.s.

Note that $W_t \sim N(0, t)$, and $(W_{t_1}, W_{t_2}, ..., W_{t_n})'$ is multivariate normal. To see the latter, we examine a bivariate case:

$$\left(\begin{array}{c} W_s \\ W_t \end{array}\right) = \left(\begin{array}{c} 1 & 0 \\ 1 & 1 \end{array}\right) \left(\begin{array}{c} W_s \\ W_t - W_s \end{array}\right).$$

It is also clear that $W_t|W_s = x \sim N(x, t - s)$. Furthermore, we can easily prove the following properties:

- Time-homogeneity $V_t = W_{t+s} - W_s$ for any fixed s is a BM independent of \mathcal{F}_s .
- Symmetry $V_t = -W_t$ is a BM.
- Scaling $V_t = cW_{t/c^2}$ is a BM.
- Time inversion $V_0 = 0, V_t = tW_{1/t}, t > 0$ is a BM.

2.2.2 Martingale

A stochastic process M_t is a martingale with respect to a filtration \mathcal{F}_t if M is adapted to \mathcal{F} and

$$\mathbb{E}(M_t | \mathcal{F}_s) = M_s \quad \text{for} \quad s < t.$$

When $\mathbb{E}(M_t|\mathcal{F}_s) \geq M_s$, we call M_t a sub-martingale. When $\mathbb{E}(M_t|\mathcal{F}_s) \leq M_s$, we call it a sup-martingale.

Remarks: (1) It is clear that Brownian motion is martingale.

(2) $W_t^2 - t$ is a martingale.

Proof Write $W_t^2 = (W_t - W_s + W_s)^2 = (W_t - W_s)^2 + W_s^2 + 2(W_t - W_s)W_s$ with s < t. Hence $\mathbb{E}(W_t^2 | \mathcal{F}_s) = (t - s) + W_s^2$. Hence $\mathbb{E}(W_t^2 - t | \mathcal{F}_s) = W_s^2 - s$.

Note that the first step in the proof is called *incrementalization*, which produces independent increments. Using the same technique, we can prove: (3) $\exp(\lambda W_t - \lambda^2 t/2)$ is a positive martingale.

Using the law of iterative expectation, we can prove: (4) For any random variable X, let $\xi_t = \mathbb{E}(X|\mathcal{F}_t)$, then $\xi = (\xi_t)$ is a martingale with respect to \mathcal{F} .

Using Jensen's inequality, we can prove: (5) $|M_t|^p$ is a submartingale if $p \ge 1$ and $\mathbb{E}|M_t|^p < \infty$.

2.2.3 Markov Process

 $(X_t, \mathcal{F}_t) \sim$ a Markov process if the distribution of X_t conditional on \mathcal{F}_s with s < t is identical to the distribution of X_t conditional on $\sigma(X_s)$. Intuitively, at time s, X_s contains all information useful for predicting X_t and all past history of X before time s is irrelevant.

The likelihood of the discrete samples of a Markov process has a nice iterative representation. Choose $t_1, t_2, ..., t_n$ arbitrarily. The likelihood of $(X_{t_1}, X_{t_1}, ..., X_{t_n})$, in general, is given by

$$f(X_{t_1}, X_{t_1}, \dots, X_{t_n}) = f(X_{t_1}) \cdot f(X_{t_2} | X_{t_1}) \cdot f(X_{t_3} | X_{t_1}, X_{t_2}) \cdots f(X_{t_n} | X_{t_1}, \dots, X_{t_{n-1}}).$$

For Markov processes, we have

$$f(X_{t_1}, X_{t_1}, \dots, X_{t_n}) = f(X_{t_1}) \cdot f(X_{t_2} | X_{t_1}) \cdot f(X_{t_3} | X_{t_2}) \cdots f(X_{t_n} | X_{t_{n-1}}).$$

So to determine the distribution of a continuous Markov process, we just need to determine the distribution of $X_t | X_s = x$ for any t and s.

Transition probability The transition probability of a Markov process (X_t) is given by

$$P_{s,t}(x,A) = \mathbb{P}\{X_t \in A | X_s = x\}.$$

The transition probability completely determines the distribution of Markov processes.

Homogenous Markov process If the transition probability of a Markov process (X_t) does not depend on the starting time,

$$\mathbb{P}\{X_{s+t} \in A | X_s = x\} = \mathbb{P}\{X_t \in A | X_0 = x\},\$$

we call it a homogenous Markov process. In this case, we have a simpler notation for transition probability,

$$P_t(x, A) = P_{s,s+t}(x, A) = \mathbb{P}\{X_t \in A | X_0 = x\}.$$

Conditional expectation operator $P_t(x, A)$ describes a distribution by assigning probability values to subsets A. We can also describe distribution by its generalized moments $\mathbb{E}f(Z)$, where $Z = X_t | X_s = x$ in our case. If we know $\mathbb{E}f(Z)$ for enough number of f, then we know the distribution of Z. In fact, we can choose f to be

$$f(\cdot) = I\{ \cdot \in A \}.$$

Then $\mathbb{E}f(Z)$ is exactly $\mathbb{P}\{Z \in A\}$,

$$\mathbb{E}f(Z) = \mathbb{E}I\{Z \in A\} = \mathbb{P}\{Z \in A\}.$$

The following notation is often used for homogenous Markov processes,

$$P_t f(x) \equiv P_t(x, f) = \mathbb{E}(f(X_t)|X_0 = x).$$
(2.1)

 P_t is linear operator on a vector space of real-valued functions:

$$P_t: f \mapsto P_t f.$$

Transition density Transition density, denoted by p(t, x, y), may be defined with respect to transition probability,

$$P_t(x,A) = \int_A p(t,x,y)dy.$$
(2.2)

Since $P_t f(x)$ is a conditional expectation, we have

$$P_t f(x) = \int_{-\infty}^{\infty} f(y) p(t, x, y) dy.$$

Example: Brownian motion. The property of independent increment ensures that the Brownian motion is a Markov process. And we have $W_{t+s}|W_s = x \sim N(x,t)$. So W_t is a homogenous Markov process. The transition density is given by

$$p(t, x, y) = \frac{1}{2\pi\sqrt{t}} \exp\left(-\frac{(y-x)^2}{2t}\right).$$

2.2.4 Chapman-Kolmogorov Equation

For any homogeneous Markov process, we have

$$P_{s+t}(x,A) = \int P_s(x,dy) P_t(y,A).$$
 (2.3)

Proof We have

$$P_{s+t}(x,A) = \mathbb{P} \left(X_{t+s} \in A | X_0 = x \right)$$

$$= \mathbb{E} \left[\mathbb{P} \left(X_{t+s} \in A | X_s \right) | X_0 = x \right]$$

$$= \mathbb{E} \left[f(X_s) | X_0 = x \right] \quad \longleftrightarrow \quad f(y) = \mathbb{P} \left(X_{t+s} \in A | X_s = y \right) = P_t(y,A)$$

$$= P_s f$$

$$= \int P_s(x,dy) f(y)$$

$$= \int P_s(x,dy) P_t(y,A).$$

In terms of the conditional expectation operator, we have for any positive measurable function f,

$$P_{s+t}f = P_s P_t f. (2.4)$$

Proof We have

$$P_{t+s}f(x) = \int P_{s+t}(x, dz)f(z)$$

=
$$\int \int P_s(x, dy)P_t(y, dz)f(z)$$

=
$$\int P_s(x, dy) \int P_t(y, dz)f(z)$$

=
$$(P_sP_tf)(x).$$

Set $f = 1_A$, this becomes the Chapman-Kolmogorov Equation (2.3).

2.3 Ito Calculus

2.3.1 Ito Integral

In this section we study the following integral:

$$\int_0^t K_s dM_s,$$

where (M_t, \mathcal{F}_t) is a continuous martingale and K is adapted to \mathcal{F} .

First we study the ordinary Lebesgue-Stieltjes integral:

$$\int_0^t f(s) dg(s).$$

For any partition $0 = t_0 < t_1 < t_2 < \cdots < t_n = t$, define

$$S = \sum_{i} f(s_i) [g(t_i) - g(t_{i-1})],$$

where $t_{i-1} \leq s_i \leq t_i$. Let $\pi_t = \max_i |t_i - t_{i-1}|$. We say that the Lebesgue-Stieltjes integral exists, if $\lim_{\pi_t \to 0} S$ exists for any $s_i \in [t_{i-1}, t_i]$.

Suppose that f is continuous and g is of bounded variation, ie,

$$\sum_{i} |g(t_i) - g(t_{i-1})| < \infty.$$

For example, if g is monotonely increasing, then it is of bounded variation. We look at

$$S1 = \sum f(s_i)[g(t_i) - g(t_{i-1})]$$

$$S2 = \sum f(t_{i-1})[g(t_i) - g(t_{i-1})].$$

And find that

$$|S_1 - S_2| \leq \left(\max_i |f(s_i) - f(t_{i-1})| \right) \left(\sum_i |g(t_i) - g(t_{i-1})| \right) \to 0.$$

So the Lebesgue-Stieltjes integral exists.

However, it is well known that a martingale M is of bounded variation if and only if M is constant. In other words, $\int_0^t K_s dM_s$ cannot be defined "path-by-path" as a Lebesgue-Stieltjes integral. Instead, we define

$$\int_{0}^{t} K_{s} dM_{s} = p \lim_{\pi_{t} \to 0} \sum_{i} K_{t_{i-1}} (M_{t_{i}} - M_{t_{i-1}}).$$
(2.5)

The choice of $K_{t_{i-1}}$ in (2.5) makes (2.5) an Ito integral. If we choose $(K_{t_{i-1}} + K_{t_i})/2$ instead of the left endpoint of each inteval, then we have Stratonovich integral. If K is bounded, measurable, and \mathcal{F}_t -adapted, then the Ito integral in (2.5) is always well defined.

Quadratic Variation For a continuous martingale M, the quadratic variation of M, denoted by [M], is defined as

$$[M]_t = \text{plim}_{\pi_t \to 0} \sum_{i=1}^n (M_{t_i} - M_{t_{i-1}})^2.$$
(2.6)

It is clear that $[M]_t$ is non-decreasing, thus of bounded variation. Thus it is integrable in the Stieltjes sense.

For any continuous process X, the first-order variation on [0, t] is captured by $\sum |X_{t_i} - X_{t_{i-1}}|$, and $\sum |X_{t_i} - X_{t_{i-1}}|^2$ captures the second order. Intuitively, for locally smooth stochastic processes, the first-order variation dominates. For locally volatile processes, the first-order variation explodes, but the second-order variation may be well defined.

Example For a Brownian motion W, we have $[W]_t = t$. To show this, partition [0, t] into n intervals of equal length $\Delta = t/n$. we have by the law of large number,

$$n\frac{1}{n}\sum_{i}(W_{i\Delta} - W_{(i-1)\Delta})^2 \to_p n\mathbb{E}(W_{i\Delta} - W_{(i-1)\Delta})^2 = n\Delta = t.$$

Quadratic Covariation Given two continuous martingales, M and N, their quadratic covariation is defined by

$$[M, N]_t = \text{plim}_{\pi_t \to 0} \sum_{i=1}^n (M_{t_i} - M_{t_{i-1}})(N_{t_i} - N_{t_{i-1}}).$$

It is straightforward to show that

$$[X+Y]_t = [X]_t + [Y]_t + 2[X,Y]_t.$$

It is also easy to show that, given a continuous martingale M and bounded-variation process A, we have

$$[M,A]_t = 0.$$

Semimartingale If X can be written as $X_t = A_t + M_t$, where (M_t) is a continuous martingale and (A_t) a continuous adapted process of finite variation, then X is called a continuous semimartingale. A constitutes trend, while M determines local variation. A continuous semimartingale X = A + M has a finite quadratic variation and $[X]_t = [M]_t$.

It is clear that Ito integral with respect to semimartingale, $\int_0^t K_s dX_s$, is well defined. We have

$$\int_0^t K_s dX_s = \int_0^t K_s dA_s + \int_0^t K_s dM_s.$$

The second item is Ito integral, and the first item is essentially a Stieltjes integral.

Properties of Ito Integral Consider $P_t = \int_0^t K_s dM_s = \text{plim}_{|\pi_t| \to 0} \sum_i K_{t_{i-1}}(M_{t_i} - M_{t_{i-1}})$, where $(M_t, \mathcal{F}_t) \sim \text{is continuous martingale}$, K_t is adapted, and $\int_0^t K_s^2 ds < \infty$ for all t. P_t has the following properties,

(a) P_t is a Martingale.

(b)
$$[P]_t = \int_0^t K_s^2 d[M]_s.$$

(c) If $P_t = \int_0^t K_s dM_s$ and $Q_t = \int_0^t H_s dN_s$, then

$$[P,Q]_t = \int_0^t K_s H_s d[M,N]_s.$$

To understand (a) intuitively, note that

$$P_{t_i} - P_{t_{i-1}} \approx K_{t_{i-1}}(M_{t_i} - M_{t_{i-1}})$$

is a martingale difference sequence. To understand (b), we write

$$[P]_{t} = \operatorname{plim} \sum (P_{t_{i}} - P_{t_{i-1}})^{2}$$

= $\operatorname{plim} \sum K_{t_{i-1}}^{2} (M_{t_{i}} - M_{t_{i-1}})^{2}$
 $\approx \operatorname{plim} \sum K_{t_{i-1}}^{2} ([M]_{t_{i}} - [M]_{t_{i-1}})^{2}$

Recall that $[M]_{t_i} = \text{plim} \sum_{k=1}^{i} (M_{t_i} - M_{t_{i-1}})^2$.

Examples (a) If $M_t = \int_0^t W_s dW_s$, then $[M]_t = \int_0^t W_s^2 ds$. (b) If $M_t = \int_0^t W_s dW_s$, then $[M, W]_t = \int_0^t W_s d[W, W]_s = \int_0^t W_s ds$.

2.3.2 Ito's Formula

Integration by parts If X and Y are two continuous semimartingales, then

$$X_t Y_t = X_0 Y_0 + \int_0^t X_s dY_s + \int_0^t Y_s dX_s + [X, Y]_t.$$
 (2.7)

In particular,

$$X_t^2 = X_0^2 + 2\int_0^t X_s dX_s + [X]_t.$$

Proof It suffices to prove the second statement. We have

$$X_{t_i}^2 - X_{t_{i-1}}^2 = 2X_{t_{i-1}}(X_{t_i} - X_{t_{i-1}}) + (X_{t_i} - X_{t_{i-1}})^2.$$

Taking sum and limit, we obtain the desired result. To prove the first statement, note that $X_t Y_t = [(X_t + Y_t)^2 - X_t^2 - Y_t^2]/2.$

In differential form, we may rewrite (2.7) as

$$d(X_tY_t) = X_t dY_t + Y_t dX_t + d[X,Y]_t.$$

Recall that for ordinary functions f(t) and g(t), we have

$$\int_{a}^{b} f(t)dg(t) = f(t)g(t)|_{a}^{b} - \int_{a}^{b} g(t)df(t).$$

Rearranging terms, we have

$$f(b)g(b) = f(a)g(a) + \int_{a}^{b} f(t)dg(t) + \int_{a}^{b} g(t)df(t).$$

Now we introduce the celebrated Ito's formula.

Ito's formula Let X be a continuous semimartingale, and $f \in C^2(\mathbb{R})$, then f(X) is a continuous semimartingale and

$$f(X_t) = f(X_0) + \int_0^t f'(X_s) dX_s + \frac{1}{2} \int_0^t f''(X_s) d[X]_s.$$
 (2.8)

In differential form, we may write the Ito's formula as

$$df(X_t) = f'(X_t)dX_t + \frac{1}{2}f''(X_t)d[X]_t.$$
(2.9)

Proof We prove by induction. Suppose $dX_t^n = nX_t^{n-1}dX_t + \frac{n(n-1)}{2}X_t^{n-2}d[X]_t$, we prove $dX_t^{n+1} = (n+1)X_t^n dX_t + \frac{n(n+1)}{2}X_t^{n-1}d[X]_t$. It is obviously true for n = 1. For arbitrary n,

$$d(X_t \cdot X_t^n) = X_t^n dX_t + X_t dX_t^n + d[X, X^n]_t$$

= $X_t^n dX_t + X_t (nX_t^{n-1} dX_t + \frac{n(n-1)}{2} X_t^{n-2} d[X]_t) + nX_t^{n-1} d[X]_t$
= $(n+1)X_t^n dX_t + \frac{n(n+1)}{2} X_t^{n-1} d[X]_t.$

So (2.9) is valid for polynomial functions. We can infer it remains true for all $f \in C^2$.

Example Since $dW_t^2 = 2W_t dW_t + dt$, $W_1^2 = \int_0^1 W_t dW_t + 1$, so we have $\int_0^1 W_t dW_t = (W_1^2 - 1)/2.$

Next we introduce the multivariate Ito's formula. Let $X = (X^1, ..., X^d)$ be a vector of continuous semimartingales and $f \in C^2(\mathbb{R}^d, \mathbb{R})$; then f(X) is a continuous semimartingale and

$$f(X_t) = f(X_0) + \sum_i \int_0^t \frac{\partial f}{\partial x_i}(X_s) dX_s^i + \frac{1}{2} \sum_{i,j} \int_0^t \frac{\partial^2 f}{\partial x_i \partial x_j}(X_s) d[X^i, X^j]_s.$$
(2.10)

In differential form, we have

$$df(X_t) = \sum_i \frac{\partial f}{\partial x_i}(X_t) dX_t^i + \frac{1}{2} \sum_{i,j} \frac{\partial^2 f}{\partial x_i \partial x_j}(X_t) d[X^i, X^j]_t.$$
(2.11)

In particular, for the bivariate case,

$$df(X_t, Y_t) = f_1(X_t, Y_t) dX_t + f_2(X_t, Y_t) dY_t + \frac{1}{2} f_{11}(X_t, Y_t) d[X]_t + f_{12}(X_t, Y_t) d[X, Y]_t + \frac{1}{2} f_{22}(X_t, Y_t) d[Y]_t$$

Furthermore, if A is of bounded variation, we have

$$df(X_t, A_t) = f_1(X_t, A_t) dX_t + f_2(X_t, A_t) dA_t + \frac{1}{2} f_{11}(X_t, A_t) d[X]_t.$$

In particular, if $dX_t = \mu_t dt + \sigma_t dW_t$, then

$$df(X_t, t) = f_1(X_t, t)dX_t + f_2(X_t, t)dt + \frac{1}{2}f_{11}(X_t, t)d[X]_t$$

= $(\mu_t f_1 + f_2 + \frac{1}{2}\sigma_t^2 f_{11})dt + \sigma_t f_1 dW_t.$

This special case often appears as "Ito's formula".

Example Consider $\xi_t = \exp(\lambda M_t - \frac{\lambda^2}{2}[M]_t) \equiv f(M_t, [M]_t)$, where M is a continuous martingale. ξ_t is called an exponential martingale. We have

$$f_1 = \lambda f$$

$$f_2 = -\frac{\lambda^2}{2} f$$

$$f_{11} = \lambda^2 f,$$

 So

$$d\exp(\lambda M_t - \frac{\lambda^2}{2}[M]_t) = \lambda \exp(\lambda M_t - \frac{\lambda^2}{2}[M]_t) dM_t,$$

or

$$\exp(\lambda M_t - \frac{\lambda^2}{2}[M]_t) = 1 + \lambda \int_0^t \exp(\lambda M_s - \frac{\lambda^2}{2}[M]_s) dM_s$$

Note that the exponential martingale is positive.

In general, if M_t is a martingale, $f(M_t)$ is not necessarily a martingale, but it is always a semimartingale. If X_t is semimartingale, the $f(X_t)$ is still semimartingale. So we say that the class of semimartingales is "invariant" under composition with C^2 -functions.

2.4 Diffusions

2.4.1 Definition and Properties

A diffusion is a continuous-time semimartingale that is characterized by the following stochastic differential equation,

$$dX_t = \mu(X_t)dt + \sigma(X_t)dW_t,$$

where $\mu(\cdot)$ is called the drift function and $\sigma(\cdot)$ is called the diffusion function. In physics, the diffusion is used to describe the movement of a particle suspended in moving liquid.

In integral form, we have

$$X_t = X_0 + \int_0^t \mu(X_s) ds + \int_0^t \sigma(X_s) dW_s.$$

Since X_t is built on a Brownian motion, it is Markov. And the fact that the functional form of $\mu(\cdot)$ and $\sigma(\cdot)$ do not change over time ensures that X_t is homogenous Markov.

Let Δ be a short time interval. On $[t, t + \Delta]$, we have

$$X_{t+\Delta} - X_t = \int_t^{t+\Delta} \mu(X_s) ds + \int_t^{t+\Delta} \sigma(X_s) dW_s.$$

It is clear that

$$\lim_{\Delta \to 0} \frac{1}{\Delta} \mathbb{E}(X_{t+\Delta} - X_t | X_t = x) = \mu(x).$$

So μ measures the rate of instantaneous changes in conditional mean. We also have,

$$\lim_{\Delta \to 0} \frac{1}{\Delta} \operatorname{var}(X_{t+\Delta} - X_t | X_t = x) = \sigma^2(x).$$

So σ^2 measures the rate of instantaneous changes in conditional volatility. It is easy to see that if μ is bounded,

$$\int_{t}^{t+\Delta} \mu(X_s) ds = O(\Delta),$$

and that if σ is bounded,

$$\int_t^{t+\Delta} \sigma(X_s) dW_s = O(\Delta^{1/2}).$$

So if we look at small intervals, the diffusion term dominates. In fact, drift term is not identifiable in small intervals. In the long run, however, the drift part dominates since π

$$\int_0^T \mu(X_s) ds = O(T),$$

while

$$\int_0^T \sigma(X_s) dWs = O(T^{1/2}).$$

Linear Drift The linear (or affine) drift function is widely used in modeling processes with mean reversion. Specifically, we may have

$$\mu(x) = \kappa(u - x),$$

where κ and u are parameters. Since $\mathbb{E}(X_{t+\Delta} - X_t) \approx \Delta \kappa (u - X_t)$. So when $\kappa > 0$, the linear drift tend to be "mean reverting", producing downward correction when $X_t > u$. However, u may or may not be the mean of the process. When $\kappa = 0$, the process is a martingale. When $\kappa < 0$, the process is unstable.

Constant-Elasticity Diffusion Many diffusion processes are endowed with the following form of diffusion function,

$$\sigma(x) = c|x|^{\rho},$$

where c and ρ are constants. From

$$\log \sigma^2(x) = \log c^2 + 2\rho \log |x|,$$

we have

$$\frac{d\log\sigma(x)}{d\log|x|} = \rho$$

This form of diffusion function is hence called constant-elasticity diffusion.

2.4.2 Useful Diffusions

In the following, we introduce a number of useful parametric diffusion models.

Brownian motion with drift

$$dX_t = \mu dt + \sigma dW_t$$

$$X_t = X_0 + \mu t + \sigma W_t$$

Transition distribution: $X_{t+\Delta} | X_t = x \sim N(x + \mu \Delta, \sigma^2 \Delta).$

Geometric Brownian motion

$$dX_t = \mu X_t dt + \sigma X_t dW_t$$

By Ito's formula,

$$d\log X_t = \frac{1}{X_t} dX_t - \frac{1}{2X_t^2} d[X]_t$$

Since $d[X]_t = \sigma^2 X_t^2 dt$,

$$d\log X_t = (\mu - \frac{1}{2}\sigma^2)dt + \sigma dW_t.$$

Ornstein-Uhlenbeck process The Ornstein-Uhlenbeck process has the following SDE representation,

$$dX_t = \kappa(\mu - X_t)dt + \sigma dW_t.$$

To derive the transition distribution, we define $Y_t = X_t - \mu$. Then

$$dY_t = -\kappa Y_t dt + \sigma dW_t.$$

$$d(\exp(\kappa t)Y_t) = \kappa \exp(\kappa t)Y_t dt + \exp(\kappa t)dY_t$$

= $\kappa \exp(\kappa t)Y_t dt + \exp(\kappa t)(-\kappa Y_t dt + \sigma dW_t)$
= $\sigma \exp(\kappa t)dW_t$

So

$$\exp(\kappa t)Y_t = Y_0 + \sigma \int_0^t \exp(\kappa s) dW_s,$$

ie,

$$Y_t = \exp(-\kappa t)Y_0 + \sigma \int_0^t \exp(-\kappa(t-s))dW_s.$$

So

$$X_t = \mu + \exp(-\kappa t)(X_0 - \mu) + \sigma \int_0^t \exp(-\kappa(t - s))dW_s.$$

Given $Y_0 = y$, what is the distribution of Y_t ?

$$Y_t = \exp(-\kappa t)Y_0 + \sigma \int_0^t \exp(-\kappa (t-s))dW_s$$

~ $N(\exp(-\kappa t)y, \sigma^2 \frac{1 - \exp(-2\kappa t)}{2\kappa}).$

Let $t \to \infty$,

$$Y_t \sim N(0, \frac{\sigma^2}{2\kappa}).$$

So if $Y_0 \sim N(0, \frac{\sigma^2}{2\kappa})$, Y_t is stationary and $Y_t \sim N(0, \frac{\sigma^2}{2\kappa})$.

Feller's squared-root process The Feller's squared-root process has the following representation,

$$dX_t = \kappa(\mu - X_t)dt + \sigma\sqrt{X_t}dW_t.$$

If $\frac{2\kappa\mu}{\sigma^2} \ge 1$, then $X_t \in [0, \infty)$. Like Ornstein-Uhlenbeck process, Feller's squared-root process is also a stationary process. And it's transition distribution is non-central Chi-square, and marginal distribution gamma. It is used by Cox, Ingersol, and Ross (CIR) to model interest rates.

Infinitestimal generator Let P_t be the condition expectation operator defined in (2.1) for a homogenous Markov process. We may understand P_0 as the identity operator, that is, $P_0 f = f$. Define the infinitestimal generator, denoted by A, as

$$Af(x) = \lim_{t \to 0} \frac{P_t f(x) - P_0 f(x)}{t} = \lim_{t \to 0} \frac{P_t f(x) - f(x)}{t}.$$
 (2.12)

If X_t follows $dX_t = \mu(X_t)dt + \sigma(X_t)dW_t$, then we have

$$f(X_t) = f(X_0) + \int_0^t (\mu f' + \frac{1}{2}\sigma^2 f'')(X_s)ds + \int_0^t (\sigma f')(X_s)dW_s.$$

Hence

$$P_t f(x) = f(x) + (\mu f' + \frac{1}{2}\sigma^2 f'')(x)t + O(t^2).$$

 So

$$Af = \lim_{t \to 0} \frac{P_t f - f}{t} = \mu f' + \frac{1}{2}\sigma^2 f''.$$

Using this infinitestimal generator, we can write $P_t f(x) = \mathbb{E}(f(X_t)|X_0 = x)$ in the form of a Taylor series expansion,

$$P_t f(x) = f(x) + A f(x)t + \frac{1}{2}A^2 f(x)t^2 + \dots + \frac{1}{j!}A^j f(x)t^j + O(t^{j+1}).$$
(2.13)

Kolmogrov forward and backward equations We have

$$\frac{d}{dt}P_t f = \lim_{s \to 0} \frac{P_{t+s}f - P_t f}{s} = \lim_{s \to 0} \frac{P_t(P_s f - f)}{s} = P_t A f \qquad (2.14)$$

$$= \lim_{s \to 0} \frac{P_s(P_t f) - (P_t f)}{s} = A P_t f \qquad (2.15)$$

(2.14) and (2.15) are Kolmogrov forward and backward equations, respectively. The above also proves that P_t and A commutes.

Using transition density (p(t, x, y)) defined in (2.2), we have

$$P_t A f(x) = \int (Af)(y)p(t,x,y)dy = \int (\mu f' + \frac{1}{2}\sigma^2 f'')(y)p(t,x,y)dy$$
$$= -\int f(y)\frac{\partial}{\partial y}(\mu(y)p(t,x,y))dy + \int f(y)\frac{\partial^2}{\partial y^2}(\frac{1}{2}\sigma^2(y)p(t,x,y))dy.$$

Since $P_t f(x) = \int f(y) p(t, x, y) dy$,

$$\frac{d}{dt}P_tf(x) = \int f(y)\frac{\partial}{\partial t}p(t,x,y).$$

So (2.14) results in

$$\frac{\partial}{\partial t}p(t,x,y) = -\frac{\partial}{\partial y}(\mu(y)p(t,x,y)) + \frac{1}{2}\frac{\partial^2}{\partial y^2}(\sigma^2(y)p(t,x,y)), \qquad (2.16)$$

which is the more common form of the Kolmogrov forward equation. Similarly,

$$\begin{aligned} AP_t f(x) &= A \int f(y) p(t, x, y) dy \\ &= (\mu(x) \frac{\partial}{\partial x} + \frac{1}{2} \sigma^2(x) \frac{\partial^2}{\partial x^2}) \int f(y) p(t, x, y) dy \\ &= \int f(y) \left(\mu(x) \frac{\partial}{\partial x} p(t, x, y) + \frac{1}{2} \sigma^2(x) \frac{\partial^2}{\partial x^2} p(t, x, y) \right) dy. \end{aligned}$$

Hence

$$\frac{\partial}{\partial t}p(t,x,y) = \mu(x)\frac{\partial}{\partial x}p(t,x,y) + \frac{1}{2}\sigma^2(x)\frac{\partial^2}{\partial x^2}p(t,x,y), \qquad (2.17)$$

which is the backward Kolmogrov equation.

2.4.3 Discrete-Time Approximation

For many diffusions, the transition distributions are very complicated. Often they do not have closed-form density functions. It is thus desirable to have approximations of transition distributions. The approximation error shall go to zero as intervals of discrete-time observations go to zero.

Euler approximation Suppose that we observe a discrete-time sequence, X_{Δ} , $X_{2\Delta}$, ..., $X_{n\Delta}$. The interval between observations is Δ . We seek an approximation of the conditional distribution of $X_{n\Delta}|X_{(n-1)\Delta}$. Let Δ be small. We have

$$\begin{aligned} X_{i\Delta} - X_{(i-1)\Delta} &= \int_{(i-1)\Delta}^{i\Delta} \mu(X_t) dt + \int_{(i-1)\Delta}^{i\Delta} \sigma(X_t) dW_t \\ &= \Delta \mu(X_{(i-1)\Delta}) + \sigma(X_{(i-1)\Delta}) (W_{i\Delta} - W_{(i-1)\Delta}) \\ &+ \int_{(i-1)\Delta}^{i\Delta} [\mu(X_t) - \mu(X_{(i-1)\Delta})] dt + \int_{(i-1)\Delta}^{i\Delta} [\sigma(X_t) - \sigma(X_{(i-1)\Delta})] dW_t \\ &\approx \Delta \mu(X_{(i-1)\Delta}) + \sigma(X_{(i-1)\Delta}) (W_{i\Delta} - W_{(i-1)\Delta}) \end{aligned}$$

This is called Euler Approximation. Under this approximation, $X_{i\Delta}|X_{(i-1)\Delta} = x \sim N(\Delta \mu(x), \Delta \sigma^2(x))$.

Milstein approximation We have something better. Consider

$$\mu(X_t) - \mu(X_{(i-1)\Delta})$$

$$= \int_{(i-1)\Delta}^t \mu'(X_s) dX_s + \frac{1}{2} \int_{(i-1)\Delta}^t \mu''(X_s) d[X]_s$$

$$= \int_{(i-1)\Delta}^t (\mu'(X_s)\mu(X_s) + \frac{1}{2}\mu''(X_s)\sigma^2(X_s)) ds + \int_{(i-1)\Delta}^t \mu'(X_s)\sigma(X_s) dW_s,$$

and

$$\sigma(X_t) - \sigma(X_{(i-1)\Delta}) = \int_{(i-1)\Delta}^t \sigma'(X_s) dX_s + \frac{1}{2} \int_{(i-1)\Delta}^t \sigma''(X_s) d[X]_s$$
$$= \int_{(i-1)\Delta}^t (\sigma'(X_s)\mu(X_s) + \frac{1}{2}\sigma''(X_s)\sigma^2(X_s)) ds + \int_{(i-1)\Delta}^t \sigma'(X_s)\sigma(X_s) dW_s.$$

And

$$\int_{(i-1)\Delta}^{i\Delta} \int_{(i-1)\Delta}^{t} (\mu\mu' + \frac{\sigma^{2}\mu''}{2}) = O(\Delta^{2})$$
$$\int_{(i-1)\Delta}^{i\Delta} \int_{(i-1)\Delta}^{t} (\sigma\mu')dW_{s}dt = O(\Delta^{3/2})$$
$$\int_{(i-1)\Delta}^{i\Delta} \int_{(i-1)\Delta}^{t} (\sigma\sigma')(X_{s})dW_{s}dW_{t} = O(\Delta) \quad (*)$$

So if we want to have an accuracy of $O(\Delta)$, the last term cannot be ignored. To have a better approximation,

$$(*) = \sigma \sigma'(X_{(i-1)\Delta}) \int_{(i-1)\Delta}^{i\Delta} \int_{(i-1)\Delta}^{t} dW_s dW_t + \int_{(i-1)\Delta}^{i\Delta} \int_{(i-1)\Delta}^{t} [(\sigma \sigma')(X_s) - (\sigma \sigma')(X_{(i-1)\Delta})] dW_s dW_t = \sigma \sigma'(X_{(i-1)\Delta}) \int_{(i-1)\Delta}^{i\Delta} (W_t - W_{(i-1)\Delta}) dW_t + o(\Delta) = \frac{1}{2} [(W_{i\Delta} - W_{(i-1)\Delta})^2 - \Delta] \sigma \sigma'(X_{(i-1)\Delta}) + o(\Delta).$$

The last equality is obtained by applying Ito's formula,

$$d(\frac{1}{2}(W_t - W_{(i-1)\Delta})^2) = (W_t - W_{(i-1)\Delta})dW_t + \frac{1}{2}dt.$$
So here's Milstein Approximation,

$$X_{i\Delta} - X_{(i-1)\Delta} = \Delta \mu(X_{(i-1)\Delta}) + \sigma(X_{(i-1)\Delta})(W_{i\Delta} - W_{(i-1)\Delta}) + \frac{1}{2}[(W_{i\Delta} - W_{(i-1)\Delta})^2 - \Delta]\sigma\sigma'(X_{(i-1)\Delta}) + o(\Delta).$$

2.4.4 Estimation*

MLE

Suppose the data generating process is parametric,

$$dX_t = \mu(X_t, \theta_0) + \sigma(X_t, \theta_0)dW_t,$$

where θ_0 is a parameter vector.

We observe X_t at evenly spaced time points, $\Delta, 2\Delta, ..., n\Delta \equiv T$. From these observations we want to estimate θ_0 .

Simple Cases When p(t, x, y) has closed-form expression (GBM, Ornstein-Uhlenbeck, CIR), we can easily form the log likelihood function as

$$L = \sum_{i=1}^{n} l(\Delta, X_{(i-1)\Delta}, X_{i\Delta}),$$

where

$$l(\Delta, X_{(i-1)\Delta}, X_{i\Delta}) = \log p(\Delta, X_{(i-1)\Delta}, X_{i\Delta}).$$

Here we may safely ignore the log likelihood function of X_0 .

Naive MLE When p(t, x, y) does not have a closed-form expression, we may apply MLE to the Euler approximation of the original diffusion,

$$X_{i\Delta} = X_{(i-1)\Delta} + \mu(X_{(i-1)\Delta}, \theta_0)\Delta + \sigma(X_{(i-1)\Delta}, \theta_0)Z_i,$$

where (Z_i) are a sequence of independent $N(0, \Delta)$ random variables.

Exact MLE We may also obtain $p(\Delta, X_{(i-1)\Delta}, X_{i\Delta})$ by solving Kolmogrov's forward or backward equation numerically. A boundary problem for the forward equation can be specified as

$$\frac{\partial}{\partial t}p(t,x,y) = -\frac{\partial}{\partial y}(\mu(y)p(t,x,y)) + \frac{1}{2}\frac{\partial^2}{\partial y^2}(\sigma^2(y)p(t,x,y)) + \frac{1}{2}\frac$$

with

$$p(0, x, y) = \delta(x - y)$$

$$p(t, x, \infty) = p(t, x, -\infty) = 0.$$

For each *i*, let $x = X_{(i-1)\Delta}$, we solve for $p(\Delta, X_{(i-1)\Delta}, X_{i\Delta})$.

Approximate MLE The idea is to construct a closed-form sequence of approximations to $p(\Delta, X_{(i-1)\Delta}, z)$: $p^{(J)}(\Delta, X_{(i-1)\Delta}, z), J = 1, 2, 3, ...$ As $J \to \infty, p^{(J)} \to p$.

We first transform X_t into a process Z_t whose transition density p_Z is close to N(0,1), making possible an expansion of p_Z around N(0,1). This involves two steps.

(1) Transform X_t into Y_t by

$$Y_t = \int^{X_t} \frac{ds}{\sigma(X_s)} =: \gamma(X_t).$$

 γ is obviously increasing and hence invertible. Using Ito's formula,

$$d\gamma(X_t) = \left(\frac{\mu(X_t)}{\sigma(X_t)} - \frac{1}{2}\sigma'(X_t)\right)dt + dW_t.$$

Hence

$$dY_t = \mu_Y(Y_t)dt + dW_t,$$

where

$$\mu_Y(Y_t) = \frac{\mu(\gamma^{-1}(Y_t))}{\sigma(\gamma^{-1}(Y_t))} - \frac{1}{2}\sigma'(\gamma^{-1}(Y_t)).$$

(2) Transform Y_t into Z_t by

$$Z_t = \Delta^{-1/2} (Y_t - y_0).$$

Now define the approximation to $p_Z(\Delta, z_0, z)$ as

$$p_Z^{(J)}(\Delta, z_0, z) = \phi(z) \sum_{j=0}^J \xi^{(j)} H_j(z),$$

where ϕ is the density function for standard normal distribution and $(H_j(z))$ are Hermite polynomials:

$$H_j(z) = e^{z^2/2} \frac{d^j}{dz^j} \left(e^{-z^2/2} \right), \ j \ge 0.$$

 $\xi^{(j)}$ satisfies

$$\xi^{(j)} = \frac{1}{j!} \int H_j(z) p_Z(\Delta, z_0, z) dz$$

= $\frac{1}{j!} \int H_j(z) \Delta^{1/2} p_Y(\Delta, y_0, \Delta^{1/2} z + y_0) dz$
= $\frac{1}{j!} \int H_j(\Delta^{-1/2} (y - y_0)) p_Y(\Delta, y_0, y) dy$
= $\frac{1}{j!} \mathbb{E} \left(H_j(\Delta^{-1/2} (Y_\Delta - y_0)) | Y_0 = y_0 \right).$ (2.18)

Note that $p_Y(\Delta, y_0, y) = \Delta^{-1/2} p_Z(\Delta, z_0, \Delta^{-1/2}(y-y_0))$. Now let $f(y) = H_j(\Delta^{-1/2}(y-y_0))$. (2.18) reduces to $P_{\Delta}f(y_0)$, which allows Taylor-type expansion,

$$P_{\Delta}f(y_0) = f(y_0) + \sum_{k=1}^{K} \frac{1}{k!} (A^k f)(y_0) \Delta^k + O(\Delta^{K+1}).$$

We choose the orders of approximation J and K. Then ξ and thus $p_Z^{(J)}(\Delta, z_0, z)$ can be explicitly calculated. We then transform $p_Z^{(J)}(\Delta, z_0, z)$ back to $p_X^{(J)}(\Delta, x_0, x)$, which is an approximation of $p_X(\Delta, x_0, x)$.

GMM

Naive GMM We have

$$dX_t = \mu(X_t, \theta_0) + \sigma(X_t, \theta_0) dW_t.$$

By Euler approximation,

$$X_{t+\Delta} \approx X_t + \mu(X_t, \theta_0) \Delta + \sigma(X_t, \theta_0) (W_{t+\Delta} - W_t).$$

Let $\varepsilon_{t+\Delta} = X_{t+\Delta} - X_t + \mu(X_t, \theta_0)\Delta$, we have

$$\mathbb{E}(\varepsilon_{t+\Delta}|X_t) = 0 \\ \mathbb{E}(\varepsilon_{t+\Delta}^2|X_t) = \sigma^2(X_t,\theta_0)\Delta$$

So we at least have four moment conditions:

$$\mathbb{E}(\varepsilon_{t+\Delta}) = 0$$
$$\mathbb{E}(\varepsilon_{t+\Delta}X_t) = 0$$
$$\mathbb{E}(\varepsilon_{t+\Delta}^2 - \sigma^2(X_t, \theta_0)\Delta) = 0$$
$$\mathbb{E}[(\varepsilon_{t+\Delta}^2 - \sigma^2(X_t, \theta_0)\Delta)X_t] = 0$$

Simulated Moment Estimation The idea is to use simulation to generate simulated moments, which are matched with sample moments.

The sample moment is simply

$$\hat{G}_n = \frac{1}{n} \sum_{i=1}^n f(X_{i\Delta}).$$

For each choice of parameter vector θ , we simulate a sequence of $X_{b\Delta}^{\theta}$, b = 1, 2, ..., B, where B is a large number. The simulated moments are thus

$$\tilde{G}(\theta) = \frac{1}{B} \sum_{b=1}^{B} f(X_{b\Delta}^{\theta}).$$

Let

$$G_n(\theta) = \tilde{G}(\theta) - \hat{G}_n.$$

The GMM estimator is given as

$$\hat{\theta}_n = \operatorname{argmin}_{\theta} G'_n(\theta) W_n G'_n(\theta),$$

where W_n is an appropriate distance matrix. See Gallant and Tauchen (1996) for more details.

The key assumption for the above strategy to work is that X_t is geometrically ergodic. Geometrical Ergodicity means that for some $\rho \in (0, 1)$, there is a probability measure P such that for any initial point x,

$$\rho^{-t} \| P_t(x, \cdot) - P \|_v \to 0 \text{ as } t \to \infty,$$

where $\|\cdot\|_{v}$ is the total variation norm defined as

$$||u||_v = \sup_A |u(A)|.$$

Exact GMM If we assume that X_t is stationary, then $\mathbb{E}f(X_t)$ does not depend on t. This leads to

$$\frac{d}{dt}\mathbb{E}f(X_t) = \lim_{\Delta \to 0} \frac{1}{\Delta} \left(\mathbb{E}f(X_{t+\Delta}) - \mathbb{E}f(X_t)\right) \\
= \mathbb{E}\left[\lim_{\Delta \to 0} \frac{1}{\Delta} \left(\mathbb{E}f(X_{t+\Delta}) - \mathbb{E}f(X_t)\right) | X_t\right] \\
= \mathbb{E}\left[\lim_{\Delta \to 0} \frac{1}{\Delta} \left(\mathbb{E}f(X_{\Delta}) - \mathbb{E}f(X_0) | X_0 = X_t\right)\right] \\
= \mathbb{E}Af(X_t) = 0$$
(2.19)

(2.19) holds for any measurable function and may serve enough number of moment conditions for GMM.

We can find more moment conditions. Define

$$P_t^* f^*(y) = \mathbb{E} \left(f^*(X_0) | X_t = y. \right),$$

where f^* is any measurable function. Obviously $P_0^* f^* = f^*$. And we define the backward infinitestimal generator

$$A^*f^* = \lim_{t \to 0} \frac{P_t^*f^* - f^*}{t}.$$

 P_t^* is the adjoint of P_t . To see this,

$$\langle f^*(X_0), P_t f(X_0) \rangle \equiv \mathbb{E} \left[f^*(X_0) \mathbb{E}(f(X_t) | X_0) \right]$$

$$= \mathbb{E} \left[f^*(X_0) f(X_t) \right]$$

$$= \mathbb{E} \left[\mathbb{E}(f(X_0) | X_t) f(X_t) \right]$$

$$= \langle P_t^* f^*(X_t), f(X_t) \rangle$$

$$= \langle P_t^* f^*(X_0), f(X_0) \rangle$$

The last equality uses the stationarity of X_t . We can also show that A^* is the adjoint of A. Then we have

$$\langle P_t A f(X_0), f^*(X_0) \rangle = \langle A P_t f(X_0), f^*(X_0) \rangle = \langle f(X_0), P_t^* A^* f^*(X_0) \rangle.$$

The inner product on the left,

$$\langle P_t A f(X_0), f^*(X_0) \rangle = \mathbb{E}[\mathbb{E}(A f(X_t) | X_0) f^*(X_0)] = \mathbb{E}[A f(X_t) f^*(X_0)].$$

The inner product on the right,

$$\langle f(X_0), P_t^* A^* f^*(X_0) \rangle = \langle f(X_t), P_t^* A^* f^*(X_t) \rangle$$

= $\mathbb{E}[f(X_t) \mathbb{E}(A^* f^*(X_0) | X_t)]$
= $\mathbb{E}[f(X_t) A^* f^*(X_0)].$

Hence

$$\mathbb{E}[Af(X_t)f^*(X_{t-\Delta}) - f(X_t)A^*f^*(X_{t-\Delta})] = 0.$$
(2.20)

(2.20) offer more choices of moment conditions for GMM. In particular, if f^* is a constant function, (2.20) reduces to (2.19).

Eigen GMM Consider the infinitestimal generator A of X_t . It is well known from the spectral theory of diffusion processes that for many diffusions, the set of eigenvalues (spectrum) Λ_{θ} for A are positive and discrete. So Λ_{θ} can be written as $(\lambda_1, \lambda_2, ..., \lambda_n, ...)$, where $0 \leq \lambda_1 < \lambda_2 < ... < \lambda_n <$

Let (λ, ϕ) be any eigen-pair of A. We have

$$A\phi = -\lambda\phi.$$

Then

$$\frac{dP_t\phi}{dt} = \lim_{\Delta \to 0} \frac{P_{t+\Delta}\phi - P_t\phi}{\Delta}$$
$$= P_tA\phi = -\lambda P_t\phi.$$

This is ordinary differential equation on $P_t\phi$. It is well known that

$$P_t \phi = e^{-\lambda t} \phi$$

Now apply Ito's formula to $e^{\lambda t}\phi(X_t)$,

$$de^{\lambda t}\phi(X_t) = \lambda e^{\lambda t}\phi(X_t)dt + e^{\lambda t}\phi'(X_t)dX_t + \frac{1}{2}e^{\lambda t}\phi''(X_t)d[X]_t$$

$$= \left(\lambda e^{\lambda t}\phi(X_t) + e^{\lambda t}\phi'(X_t)\mu(X_t) + \frac{1}{2}e^{\lambda t}\phi''(X_t)\sigma^2(X_t)\right)dt$$

$$+ e^{\lambda t}\phi'(X_t)\sigma(X_t)dW_t$$

$$= e^{\lambda t}(\lambda\phi(X_t) + A\phi(X_t))dt + e^{\lambda t}\phi'(X_t)\sigma(X_t)dW_t$$

$$= e^{\lambda t}\phi'(X_t)\sigma(X_t)dW_t.$$

Hence,

$$\phi(X_t) = e^{-\lambda t}\phi(X_0) + \int_0^t e^{-\lambda(t-s)}\phi'(X_s)\sigma(X_s)dW_s.$$

 So

$$\mathbb{E}(\phi(X_{t+\Delta})|X_t) = e^{-\lambda\Delta}\phi(X_t),$$

which leads to desired moment condition,

$$E\left[\left(\phi(X_{t+\Delta}) - e^{\lambda\Delta}\phi(X_t)\right)g(X_t)\right] = 0, \qquad (2.21)$$

where g can be any measurable function.

Nonparametric Estimation

The methodology of MLE and GMM presupposes correct parameterization of the diffusion models (or equivalently the infinitestimal generator). Any misspecification leads to inconsistency of those estimators. The problem of parameterization can be avoided by the use of nonparametric diffusion models, the time-homogeneous version of which is given as,

$$dX_t = \mu(X_t)dt + \sigma(X_t)dW_t.$$

Ignoring terms in (2.13) that are of smaller order than $O(\Delta)$, we obtain

$$\mathbb{E}[f(X_{t+\Delta})|X_t = x] = f(x) + Af(x)\Delta + O(\Delta^2),$$

where $Af(x) = \mu(x)f'(x) + \frac{1}{2}\sigma^2(x)f''(x)$. Then we have

$$Af(x) = \frac{1}{\Delta} \mathbb{E}[(f(X_{t+\Delta}) - f(X_t))|X_t = x] + O(\Delta).$$

Let f be such that f(x) = x. Then we have

$$Af(x) = \mu(x) = \frac{1}{\Delta} \mathbb{E}[(X_{t+\Delta} - X_t) | X_t = x] + O(\Delta).$$

Let $f(x) = (x - X_t)^2$. We have $Af(x) = 2\mu(x)(x - X_t) + \sigma^2(x)$, and

$$\sigma^2(x) = \frac{1}{\Delta} \mathbb{E}[(X_{t+\Delta} - X_t)^2 | X_t = x] + O(\Delta).$$

We can estimate $\mathbb{E}[(X_{t+\Delta} - X_t)|X_t = x]$ and $\mathbb{E}[(X_{t+\Delta} - X_t)^2|X_t = x]$ using Nadaraya-Watson kernel estimator.

$$\hat{\mu}(x) = \frac{\sum_{i=1}^{n} K(\frac{X_{(i-1)\Delta} - x}{h})(X_{i\Delta} - X_{(i-1)\Delta})}{\Delta \sum_{i=1}^{n} K(\frac{X_{(i-1)\Delta} - x}{h})}$$
$$\hat{\sigma}^{2}(x) = \frac{\sum_{i=1}^{n} K(\frac{X_{(i-1)\Delta} - x}{h})(X_{i\Delta} - X_{(i-1)\Delta})^{2}}{\Delta \sum_{i=1}^{n} K(\frac{X_{(i-1)\Delta} - x}{h})}.$$

The Nadaraya-Watson estimators are consistent. To see this, first note that

$$\frac{1}{nh}\sum_{i=1}^{n} K\left(\frac{X_{(i-1)\Delta}}{h}\right) \approx \frac{1}{Th}\int_{0}^{T} K\left(\frac{X_{s}-x}{h}\right) ds$$
$$= \frac{1}{Th}\int_{-\infty}^{\infty} K\left(\frac{s-x}{h}\right) L(T,s) ds$$

$$= \frac{1}{T} \int_{-\infty}^{\infty} K(s) L(T, x + hs) ds$$

$$\rightarrow \frac{1}{T} \int_{-\infty}^{\infty} K(s) L(T, x) ds$$

$$= \frac{1}{T} L(T, x),$$

where L(T, x) denotes the local time of X_t , and the derivation uses the Occupation Time Formula,

$$\int_0^t f(X_s) ds = \int_{-\infty}^\infty f(x) L(T, x) dx.$$

And

$$\frac{1}{nh} \sum_{i=1}^{n} K\left(\frac{X_{(i-1)\Delta} - x}{h}\right) \left(\frac{X_{i\Delta} - X_{(i-1)\Delta}}{\Delta}\right)$$

$$\approx \quad \frac{1}{Th} \sum K\left(\frac{X_{(i-1)\Delta} - x}{h}\right) \mu(X_{(i-1)\Delta})\Delta$$

$$\approx \quad \frac{1}{Th} \int_{0}^{t} K\left(\frac{X_s - x}{h}\right) \mu(X_s) ds$$

$$= \quad \frac{1}{Th} \int_{-\infty}^{\infty} K\left(\frac{s - x}{h}\right) \mu(s) L(T, s) ds$$

$$\rightarrow \quad \frac{1}{T} \mu(x) L(T, x).$$

We may obtain better precision by keeping more terms in (2.13). For example, we have

$$\mathbb{E}[f(X_{t+\Delta})|X_t = x] = f(x) + Af(x)\Delta + \frac{1}{2}A^2f(x)\Delta^2 + O(\Delta^3), \qquad (2.22)$$

and

$$\mathbb{E}[f(X_{t+2\Delta})|X_t = x] = f(x) + Af(x)2\Delta + \frac{1}{2}A^2f(x)4\Delta^2 + O(\Delta^3).$$
(2.23)

4(2.22)-(2.23) would give us

$$Af(x) = \frac{1}{2\Delta} \{ 4\mathbb{E}[f(X_{t+\Delta}) - f(X_t) | X_t = x] - \mathbb{E}[f(X_{t+2\Delta}) - f(X_t) | X_t = x] \} + O(\Delta^2).$$

More precise estimators of μ and σ^2 (in the order of $\Delta^2)$ then follow.

Semiparametric Estimation

We consider stationary diffusions satisfying either

$$dX_t = \mu(X_t)dt + \sigma(X_t, \theta)dW_t, \qquad (2.24)$$

or

$$dX_t = \mu(X_t, \theta)dt + \sigma(X_t)dW_t.$$
(2.25)

From Kolmogrov Forward Equation, we have

$$\begin{split} \frac{\partial}{\partial \Delta} p(\Delta, x, y) &= -\frac{\partial}{\partial y} (\mu(y) p(\Delta, x, y)) \\ &+ \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2(y) p(\Delta, x, y)). \end{split}$$

By stationarity, the density of marginal distribution is time-invariant, ie,

$$\begin{aligned} \frac{\partial}{\partial \Delta} \pi(y) &= \frac{\partial}{\partial \Delta} \int_0^\infty p(\Delta, x, y) \pi(x) dx \\ &= \int_0^\infty \frac{\partial}{\partial \Delta} p(\Delta, x, y) \pi(x) dx \\ &= \int_0^\infty \left(-\frac{\partial}{\partial y} (\mu(y) p(\Delta, x, y)) + \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2(y) p(\Delta, x, y)) \right) \pi(x) dx \\ &= -\frac{\partial}{\partial y} \left(\mu(y) \left(\int_0^\infty p(\Delta, x, y) \pi(x) dx \right) \right) \\ &\quad + \frac{1}{2} \frac{\partial^2}{\partial y^2} \left(\sigma^2(X_{t+\Delta}) \left(\int_0^\infty p(\Delta, x, y) \pi(x) dx \right) \right) \\ &= -\frac{\partial}{\partial y} \left(\mu(y) \pi(y) \right) + \frac{1}{2} \frac{\partial^2}{\partial y^2} \left(\sigma^2(y) \pi(y) \right) \\ &= 0. \end{aligned}$$

Hence

$$\frac{d^2}{x^2}(\sigma^2(x)\pi(x)) = 2\frac{d}{dx}(\mu(x)\pi(x)).$$
(2.26)

Suppose $\pi(0) = 0$, we have

$$\mu(x) = \frac{1}{2\pi(x)} \frac{d}{dx} (\sigma^2(x)\pi(x)), \qquad (2.27)$$

and

$$\sigma^{2}(x) = \frac{2}{\pi(x)} \int_{0}^{x} \mu(u)\pi(u)du.$$
(2.28)

For model (2.24), in which we have prior knowledge on the structure of σ^2 , we may parametrically estimate σ^2 and nonparametrically estimate μ using (2.27). For model (2.25), we may similarly estimate σ^2 nonparametrically using (2.28), with prior knowledge of μ .

Chapter 3

No-Arbitrage Pricing in Continuous Time

3.1 Basic Setup

Consider a financial market with an interest-paying *money account* and a *stock*.

Money Account The interest rate may be fixed, time-varying, or even statecontingent. Let M_0 be the initial deposit and M_t be the cash value of the account at time t. We may represent M_t in stochastic differential equation form as follows,

• r fixed

$$dM_t = rM_0e^{rt}dt = rM_tdt, \quad M_t = M_0e^{rt}.$$

• r time-varying, r_t

$$dM_t = r_t M_t dt, \quad M_t = M_0 e^{\int_0^t r_s ds}.$$

• r state-contingent, $r(X_t)$

$$dM_t = r(X_t)M_t dt, \quad M_t = M_0 e^{\int_0^t r(X_s)ds}$$

4

Stock Let S_t be stock price at time t that follows an Ito process,

$$dS_t = \mu(S_t)dt + \sigma(S_t)dW_t$$

The use of continuous diffusion process implicitly asserts that there is no "surprise" in the market.

One special case is when $\mu(S_t) = 0$ and $\sigma(S_t) = 1$. S_t then becomes a Brownian motion. Louis Bachelier, the pioneer of mathematical finance, used Brownian motion in describing the fluctuation in financial markets. Another well-known special case is the geometric Brownian motion, which is used by Black and Scholes (1973) to price European options.

If we assume $S_t > 0$ a.s., we may also represent S_t in the geometric form,

$$dS_t = \mu(S_t)S_t dt + \sigma(S_t)S_t dW_t.$$

Note that, throughout the text, we use homogeneous diffusions to model stock prices. More generally, at the cost of technical complication, we may also use heterogenous diffusions.

Multivariate Case S_t can be a $N \times 1$ price vector describing the prices of N securities. Accordingly, W may be a $d \times 1$ vector of independent Brownian motions, each of which represents a source of new information or innovation. In such case, $\mu \in \mathbb{R}^N$, $\sigma \in \mathbb{R}^{N \times d}$.

Portfolio/trading strategy Portfolio or trading strategy (h_t) is an adapted vector process:

$$h_t = \left(\begin{array}{c} a_t \\ b_t \end{array}\right),$$

where a_t is the holding of money account, and b_t the holding of stocks.

Trading Gain

$$G_t = \int_0^t b_s dS_s.$$

We usually impose the following integrability condition:

$$\int_0^t b_s^2 ds < \infty \text{a.s. } \forall t$$

If S_t is a martingale (e.g., BM), then we know G_t is also a martingale.

 $G_t = \int_0^t b_s dS_s$ is often called "gains process". To see this, imagine an investor who makes decisions in discrete time: $0 = t_0 < t_1 < \cdots < t_n = T$. Let b_{t_i} be the number of stocks the investor holds over the period $[t_i, t_{i+1})$. Then the gains process is described by the following stochastic difference equation:

$$G_0 = 0, \ G(t_{i+1}) - G(t_i) = b_{t_i}(S(t_{i+1}) - S(t_i)).$$

Or in summation form,

$$G(t_{i+1}) = G(0) + \sum_{j=0}^{i} b_{t_j}(S(t_{j+1}) - S(t_j)).$$

Note that as in the definition of Ito integral, b_{t_i} must be \mathcal{F}_{t_i} -measurable, meaning that the investor cannot anticipate the future (exclusion of inside trading).

Let $X_t = (M_t, S_t)'$, and let H_t be the value of the portfolio (h_t) . Then

$$H_t = h_t \cdot X_t = a_t M_t + b_t \cdot S_t.$$

Definition 3.1.1 (Self-financing) (h_t) is self-financing iff

$$dH_t = h_t \cdot dX_t = a_t dM_t + b_t \cdot dS_t.$$

Definition 3.1.2 (Arbitrage) Let (h_t) be a self-financing portfolio and (H_t) be its value, an arbitrage portfolio is one such that

$$H_0 = 0$$
, and $H_T > 0$ a.s.

Lemma 3.1.3 If there is no arbitrage opportunities, and if (h_t) is self-financing and $dH_t = v_t H_t dt$, then $v_t = r_t$, the risk-free short rate.

In other words, there is only one risk-free short rate.

Numeraire A numeraire is a strictly positive Ito process used for the "units" of pricing. If there is a riskfree rate r_t , the typical numeraire is the reciprocal of the price of riskfree zero-coupon bond, $Y_t = M_t^{-1} = \exp(-\int_0^t r_s ds)$. We denote the numeraire-deflated price process of X_t by Y_t as X^Y , $X_t^Y = X_t Y_t$. For example, if $X_t = (M_t, S_t)'$ and $Y_t = M_t^{-1}$, then $X_t^Y = (1, S_t/M_t)$.

Theorem 3.1.4 (Numeraire Invariance Theorem) Suppose Y is a numeraire. Then a trading strategy (h_t) is self-financing w.r.t. X iff (h_t) is self-financing w.r.t. X^Y . **Proof** Let $H_t = h_t \cdot X_t$, and $H_t^Y = H_t Y_t$. If $dH_t = h_t \cdot dX_t$, then

$$dH_t^Y = Y_t dH_t + H_t dY_t + d[H, Y]_t$$

= $Y_t h_t \cdot dX_t + (h_t \cdot X_t) dY_t + h_t \cdot d[X, Y]_t$
= $h_t \cdot (Y_t dX_t + X_t dY_t + d[X, Y]_t)$
= $h_t \cdot dX_t^Y$.

So (h_t) is self-financing w.r.t. X^Y . The reverse is also true.

It follows that h is an arbitrage w.r.t. X iff it is an arbitrage w.r.t. X^{Y} . All this says that renormalization of security prices by a numeraire does not have economic effects.

3.2 The Black-Scholes Model

In a market of (M_t, S_t) we price an European call option using no-arbitrage argument. We specify

$$\begin{cases} dM_t = rM_t dt \\ dS_t = \mu S_t dt + \sigma S_t dW_t \end{cases}$$

Recall that $C_T = \max(S_T - K, 0)$. In general, we may use the same argument to price any European option with the final payoff $g(S_T)$, where g is a known function.

We assume that the option price C_t can be characterized by a function of the underlying stock price S_t and time t, $C_t = F(S_t, t)$. And we assume $F \in C^{2,1}(\mathbb{R} \times [0,T))$, ie, $F_1 \equiv \frac{\partial F(x,t)}{\partial x}$, $F_2 \equiv \frac{\partial F(x,t)}{\partial t}$, and $F_{11} \equiv \frac{\partial^2 F(x,t)}{\partial x^2}$ exist and are continuous. In the options market, F_1 , F_2 , F_{11} are called delta, theta, gamma, respectively.

Using Ito's formula, we have

$$dC_t = F_2(S_t, t)dt + F_1(S_t, t)dS_t + \frac{1}{2}F_{11}(S_t, t)d[S]_t$$

= $(F_2(S_t, t) + \mu S_t F_1(S_t, t) + \frac{1}{2}F_{11}(S_t, t)\sigma^2 S_t^2)dt + \sigma S_t F_1(S_t, t)dW_t$

The pricing strategy is to replicate C_t using a self-financing portfolio of M_t and S_t . The price of the replication portfolio must then be the price of the option, if no arbitrage is allowed.

We have $H_t = a_t M_t + b_t S_t$. Since h_t is self-financing,

$$dH_t = a_t dM_t + b_t dS_t$$

= $a_t r M_t dt + b_t (\mu S_t dt + \sigma S_t dW_t)$
= $(a_t r M_t + b_t \mu S_t) dt + \sigma b_t S_t dW_t$

By the unique decomposition property of diffusion processes, since $C_t = H_t$ a.s. for all t, we have

$$b_t = F_1(S_t, t) a_t = \frac{F(S_t, t) - F_1(S_t, t)S_t}{M_t},$$

and by the equality of drift terms between C_t and H_t ,

$$F_2(S_t, t) + rS_t F_1(S_t, t) + \frac{1}{2}\sigma^2 S_t^2 F_{11}(S_t, t) - rF(S_t, t) = 0.$$
(3.1)

For (3.1) to hold, F must be the solution to the following partial differential equation (PDE):

$$F_2(x,t) + rxF_1(x,t) + \frac{1}{2}\sigma^2 x^2 F_{11}(x,t) - rF(x,t) = 0$$
(3.2)

with the boundary condition

$$F(x,T) = \max(x - K, 0).$$
(3.3)

We can check that the Black-Scholes Option Pricing Formula solves the PDE (3.2) and (3.3). The formula is as follows,

$$F(x,t) = x\Phi(z) - \exp(-r(T-t))K\Phi(z - \sigma\sqrt{T-t}), \qquad (3.4)$$

with

$$z = \frac{\log(x/K) + (r + \sigma^2/2)(T - t)}{\sigma\sqrt{T - t}},$$
(3.5)

where Φ is the cdf of standard normal distribution.

A byproduct of this derivation is a popular dynamic hedging strategy called "delta hedging". Consider a bank that has sold an European call option and now it wants to hedge its position. All it has to do is to maintain opposite positions of $h_t = (a_t, b_t)$. In fact $b_t = F_1(S_t, t)$ is called the "delta" of the option in practice.

The General Case

Now assume that (M_t, S_t) are such that

$$dM_t = r_t M_t dt$$

$$dS_t = \mu(S_t) dt + \sigma(S_t) dW_t$$

We may mimic the argument in previous section and show that under the no-arbitrage condition, the price process for an European call option $F(S_t, t)$ must satisfy:

$$F_2(x,t) + r_t x F_1(x,t) + \frac{1}{2}\sigma(x)^2 F_{11}(x,t) - r_t F(x,t) = 0$$
(3.6)

with the boundary condition

$$F(x,T) = \max(x - K, 0).$$

Note that for general European option with payoff $g(S_T)$, the price process still satisfy (3.6) with following boundary condition

$$F(x,T) = g(x). \tag{3.7}$$

3.3 The Feynman-Kac Solution

Constant Riskfree Rate

Consider the following boundary value problem:

$$F_2(x,t) + rxF_1(x,t) + \frac{1}{2}\sigma^2(x)F_{11}(x,t) - rF(x,t) = 0, \qquad (3.8)$$

with

$$F(x,T) = g(x).$$

This problem differs from (3.6) only in the form of r, which is a constant here.

Construct an Ito process Z such that $Z_t = x$

$$dZ_s = rZ_s ds + \sigma(Z_s) dW_s, \ s > t.$$

By Ito's formula,

$$d(e^{-rs}F(Z_s,s))$$

$$= [-re^{-rs}F(Z_s,s) + e^{-rs}F_2(Z_s,s)]ds + e^{-rs}F_1(Z_s,s)dZ_s + \frac{1}{2}F_{11}(Z_s,s)d[Z]_s$$

$$= e^{-rs}[-rF + F_2 + rZ_sF_1 + \frac{1}{2}\sigma^2(Z_s)F_{11}]ds + e^{-rs}\sigma(Z_s)F_1(Z_s,s)dW_s$$

If F(x,t) satisfies (3.8), then the term in bracket is zero. Hence $e^{-rs}F(Z_s,s)$ is martingale. So

$$e^{-rT}F(Z_T,T) = e^{-rt}F(Z_t,t) + \int_t^T e^{-rs}\sigma(Z_s)F_1(Z_s,s)dW_s.$$

Taking conditional expectation given $Z_t = x$ gives:

$$\mathbb{E}(e^{-rT}F(Z_T,T)|Z_t=x) \equiv \mathbb{E}(e^{-rT}g(Z_T)|Z_t=x) = e^{-rt}F(x,t).$$

Hence

$$F(x,t) = \mathbb{E}(e^{-r(T-t)}g(Z_T)|Z_t = x)$$

The message is that we can solve certain PDE's by calculating a conditional expectation of an imagined random variable $g(Z_T)$.

Stochastic Riskfree Rate

Now we work to solve (3.6) which is reproduced below,

$$F_2(x,t) + r_t x F_1(x,t) + \frac{1}{2}\sigma(x)^2 F_{11}(x,t) - r(x)F(x,t) = 0$$

with

$$F(x,T) = g(x).$$

It can be shown that if we construct Z such that $Z_t = x$ and

$$dZ_s = r_s Z_s ds + \sigma(Z_s) dW_s, \ s > t, \tag{3.9}$$

then

$$F(x,t) = \mathbb{E}\left\{\exp\left[-\int_{t}^{T} r_{s} ds\right] g(Z_{T}) | Z_{t} = x\right\}.$$
(3.10)

In particular, the Black-Scholes option price is given by

$$F(x,t) = \mathbb{E}\left\{e^{-r(T-t)}g(Z_T)|Z_t = x\right\},\qquad(3.11)$$

where Z is such that $Z_t = x$ and

$$dZ_s = rZ_s ds + \sigma Z_s dW_s, \ s > t. \tag{3.12}$$

The expectation in the Feynman-Kac solution (3.10) is taken with respect to objective probability on an imagined r.v. $g(Z_T)$. We can also represent the solution as an expectation taken with respect to an imagined probability (riskneutral probability) on a "real" r.v., for example, $g(S_T)$. This will be explored in the next section.

Calculation of Black-Scholes Formula

Now we derive Black-Scholes formula from (3.11) and (3.12). We have

$$d\log Z_s = (r - \sigma^2/2)dt + \sigma dW_t,$$

which implies

$$\log Z_T - \log Z_t = (r - \sigma^2/2)(T - t) + \sigma(W_T - W_t).$$

This is,

$$Z_T = Z_t e^{(r - \sigma^2/2)(T - t) + \sigma(W_T - W_t)}.$$

Hence

$$Z_T|_{Z_t=x} = d x e^{(r-\sigma^2/2)(T-t)+\sigma\sqrt{T-t}Z}, \ Z \sim N(0,1).$$

 So

$$F(x,t) = \mathbb{E} \left\{ e^{-r(T-t)} g(Z_T) | Z_t \right\} \\ = e^{-r(T-t)} \int_{-\infty}^{\infty} \max(x e^{(r-\sigma^2/2)(T-t) + \sigma\sqrt{T-t}z} - K, 0) \phi(z) dz$$

Some calculations yield the Black-Scholes formula in (3.4).

Feynman-Kac in Multivariate Case

Now we consider the market of a money account and multiple stocks containing d dimensions of risk. We have

$$dM_t = r_t M_t dt$$

$$dS_t = \mu(S_t) dt + \sigma(S_t) dW_t,$$

where $W_t \in \mathbb{R}^d, S_t \in \mathbb{R}^N, \mu : \mathbb{R}^N \times [0, \infty) \to \mathbb{R}^N, \sigma : \mathbb{R}^N \times [0, \infty) \to \mathbb{R}^{N \times d}.$

The option price $F(x,t): \mathbb{R}^N \times [0,\infty) \to \mathbb{R}$ solves the following boundary value pde:

$$F_1(x,t)r_t x + F_2(x,t) + \frac{1}{2} \operatorname{tr}\left[(\sigma\sigma')(x)F_{11}(x,t)\right] - r_t F(x,t) = 0$$
(3.13)

with F(x,T) = g(x). The solution of (3.13) is of the same form as (3.10).

3.4 Risk-Neutral Pricing

We have shown that in the market of M_t and S_t such that

$$dM_t = r_t M_t dt$$

$$dS_t = \mu(S_t) dt + \sigma(S_t) dW_t$$

The price of a general derivative can be represented as

$$F(x,t) = \mathbb{E}\left\{\exp\left[-\int_{t}^{T} r_{s} ds\right] g(Z_{T}) | Z_{t} = x\right\},\$$

where Z is an imagined Ito process such that $Z_s = x$ for $s \leq t$ and

$$dZ_s = r_s Z_s ds + \sigma(Z_s) dW_s, \ s > t.$$

In this section we show that there exists a probability measure $\tilde{\mathbb{P}}$ and a $\tilde{\mathbb{P}}\text{-BM}$ \tilde{W} such that

$$dS_t = r_t S_t dt + \sigma(S_t) d\tilde{W}_t.$$

So the price function of a general European option $(C_T = g(S_T))$ can be written as

$$F(x,t) = \tilde{\mathbb{E}}\left\{\exp\left[-\int_{t}^{T} r_{s} ds\right] g(S_{T}) | S_{t} = x\right\}.$$

Put it differently,

$$\frac{C_t}{M_t} = \tilde{\mathbb{E}}_t \left(\frac{C_T}{M_T} \right)$$

where M_t is the money account and acts as a numeraire. In other words, (C_t/M_t) is a martingale under $\tilde{\mathbb{P}}$. So $\tilde{\mathbb{P}}$ is sometimes called "martingale probability measure". And since the price of an asset is the expectation of its payoff C_T/M_T taken with respect to a probability measure, we call this measure "risk-neutral" measure or probability.

Change of Measure

Suppose we have a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and a nonnegative r.v. ξ that satisfies $\mathbb{E}\xi = 1$, then we may define a new probability measure as follows,

$$\tilde{\mathbb{P}}(A) = \int_A \xi(\omega) dP(\omega) \text{ for all } A \in \mathcal{F}.$$

We may check that $\tilde{\mathbb{P}}(A) \geq 0$ for all $A \in \mathcal{F}$ and $\tilde{\mathbb{P}}(\Omega) = 1$.

If $\xi > 0$ a.s., then $\tilde{\mathbb{P}}$ is called an "equivalent probability measure" of \mathbb{P} , ie, for any set A, $\tilde{\mathbb{P}}(A) = 0$ iff $\mathbb{P}(A) = 0$. ξ is called the Radon-Nikodym density of $\tilde{\mathbb{P}}$ w.r.t. \mathbb{P} . In differential form, we may write

$$d\tilde{\mathbb{P}} = \xi d\mathbb{P}.$$

For any r.v. X, it is easy to show that

$$\tilde{\mathbb{E}}X = \mathbb{E}\xi X,$$

and that

$$\mathbb{E}X = \tilde{\mathbb{E}}\frac{X}{\xi}.$$

Example Let X be standard normal, $\xi = \exp(\lambda X - \lambda^2/2)$, and $\tilde{\mathbb{P}}$ be defined as above. For any function f, we have

$$\mathbb{E}f(X) = \mathbb{E}\xi f(X)$$

= $\int \exp(\lambda x - \lambda^2/2) \frac{1}{\sqrt{2\pi}} \exp(-x^2/2) f(x) dx$
= $\frac{1}{\sqrt{2\pi}} \int \exp(-(x-\lambda)^2/2) f(x) dx.$

So under $\tilde{\mathbb{P}}$, $X \sim N(\lambda, 1)$. In other words, this particular change of measure shifts X by a constant, without changing the variance.

Martingale Equivalent Measure

Recall that there is no arbitrage opportunity if and only if there exists a risk-neutral probability measure $\tilde{\mathbb{P}}$ such that asset prices satisfy,

$$p_t = e^{-r(T-t)} \tilde{\mathbb{E}}_t x_T,$$

where r is risk-free rate. Noting that $e^{-r(T-t)} = M_t/M_T$, we re-write the above equation,

$$\frac{p_t}{M_t} = \tilde{\mathbb{E}}_t \frac{x_T}{M_T}.$$

 $\tilde{\mathbb{P}}$ is said to be the *martingale equivalent measure* of \mathbb{P} for the process p_t/M_t , since (p_t/M_t) is a martingale under $\tilde{\mathbb{P}}$.

Obviously, if a price process X_t (e.g., p_t/M_t) admits an equivalent martingale measure, then there is no arbitrage. To see this, note that for any admissible trading strategy, $\tilde{\mathbb{E}}(\int_0^t h_s dX_s) = 0$. Hence, the self-financing condition $h_t X_t = h_0 X_0 + \int_0^t h_s dX_s$ implies

$$h_0 X_0 = \tilde{\mathbb{E}}_0(h_t X_t)$$

Thus, if $h_t X_t > 0$, then $h_0 X_0 > 0$.

Density Process. Let $\xi > 0$ a.s. and $\mathbb{E}\xi = 1$ and $d\tilde{\mathbb{P}} = \xi d\mathbb{P}$. We define $\xi_t = \mathbb{E}(\xi | \mathcal{F}_t)$, which is called the density process for $\tilde{\mathbb{P}}$ with respect to \mathbb{P} . Obviously ξ_t is a positive martingale. If X_t is \mathcal{F}_t -measurable, then we have

$$\begin{aligned}
\mathbb{E}(\xi_t X_t) &= \mathbb{E}(\xi X_t) \\
\tilde{\mathbb{E}} X_t &= \mathbb{E} \xi_t X_t \\
\tilde{\mathbb{E}} \xi_t^{-1} X_t &= \mathbb{E} X_t.
\end{aligned}$$

The first statement is an immediate consequence of the law of iterative expectation, and the second statement is due to

$$\tilde{\mathbb{E}}X_t = \mathbb{E}\xi X_t = \mathbb{E}[\mathbb{E}\xi X_t | \mathcal{F}_t] = \mathbb{E}[\mathbb{E}[\xi | \mathcal{F}_t] X_t] = \mathbb{E}\xi_t X_t.$$

The third statement is similarly established. We also have

Bayes Rule.

$$\dot{\mathbb{E}}_s X_t = \xi_s^{-1} \mathbb{E}_s(\xi_t X_t), \ s < t, \tag{3.14}$$

where X_t is \mathcal{F}_t -measurable and $\tilde{\mathbb{E}}|X_t| < \infty$.

Proof For any $A \in \mathcal{F}_s$, since $d\mathbb{P} = \xi^{-1} d\tilde{\mathbb{P}}$,

$$\tilde{\mathbb{E}}\left[I_A \xi_s^{-1} \mathbb{E}_s(\xi_t X_t)\right] = \mathbb{E}\left[I_A \mathbb{E}_s(\xi_t X_t)\right] \\
= \mathbb{E}\left[I_A \xi_t X_t\right] \\
= \tilde{\mathbb{E}}\left[I_A X_t\right] \\
= \tilde{\mathbb{E}}\left[I_A \tilde{\mathbb{E}}_s X_t\right].$$

Since it is true for every A, the proof is complete.

As a result, we have the following lemma,

Lemma: If $(X_t\xi_t)$ is \mathbb{P} -martingale, then (X_t) is $\tilde{\mathbb{P}}$ -martingale.

Girsanov Theorem

Girsanov Theorem. If (X_t) is \mathbb{P} -martingale, and (\tilde{X}_t) is defined as $d\tilde{X}_t = dX_t - \xi_t^{-1}d[X,\xi]_t$, then (\tilde{X}_t) is \mathbb{P} -martingale. In other words, \mathbb{P} is the martingale equivalent measure for \tilde{X} .

Proof It suffices to show that $(\tilde{X}_t \xi_t)$ is \mathbb{P} -martingale.

$$d(\tilde{X}_t\xi_t) = \tilde{X}_t d\xi_t + \xi_t d\tilde{X}_t + d[\tilde{X},\xi]_t$$

= $\tilde{X}_t d\xi_t + \xi_t dX_t.$

Note that $[\tilde{X},\xi]_t = [X,\xi]_t$, since \tilde{X}_t and X_t differ only by a bounded-variation process $\int_0^t \xi_s^{-1} d[X,\xi]_s$.

If $X_t = W_t$, where W_t is a Brownian motion, then $d\tilde{W}_t = dW_t - \xi_t^{-1}d[W,\xi]_t$ is a Brownian motion under $\tilde{\mathbb{P}}$.

A Useful Special Case. Let η_t be adapted to \mathcal{F}_t and let $\xi_t = \exp\left(-\int_0^t \eta_s dW_s - \frac{1}{2}\int_0^t \eta_s^2 ds\right)$. If W_t is \mathbb{P} -BM, then $\tilde{W}_t = W_t + \int_0^t \eta_s ds$ is BM under $\tilde{\mathbb{P}}$.

Proof Let $L_t = -\int_0^t \eta_s dW_s$, then $\xi_t = \exp\left(L_t - \frac{1}{2}[L]_t\right)$. Then we have

$$d\log \xi_t = d(L_t - \frac{1}{2}[L]_t) = dL_t - \frac{1}{2}d[L]_t$$

and

$$d\log \xi_t = \xi_t^{-1} d\xi_t - \frac{1}{2} \frac{1}{\xi_t^2} d[\xi]_t.$$

By unique decomposition of semimartingale, we have

$$dL_t = \xi_t^{-1} d\xi_t.$$

Hence

$$\xi_t^{-1} d[W, \xi]_t = d[W, L]_t = -\eta_t dt.$$

To show that \tilde{W}_t is $\tilde{\mathbb{P}}$ -BM, we note that $[\tilde{W}]_t = t$.

Note that in the above notations, ξ_t is called the stochastic exponential of L_t .

Pricing of Contingent Claims

Using Girsanov Theorem, we can find the risk-neutral measure under which the price (in numeraire) of a contingent claim is a martingale process. Suppose we have a money account with riskfree rate r_t and a contingent claim with terminal value X_T realized at time T. Then we can find probability measure \tilde{P} under which X_t/M_t is a martingale, where X_t is the market value of the contingent claim at time t. That is, $\tilde{\mathbb{E}}_t(X_T/M_T) = X_t/M_t$, where the expectation \tilde{E} is taken with respect to \tilde{P} . Hence we have a risk-neutral pricing formula,

$$X_t = M_t \tilde{\mathbb{E}}_t \left(\frac{X_T}{M_T} \right) = \tilde{\mathbb{E}}_t \left(\exp\left(-\int_t^T r_s ds \right) X_T \right).$$

Example: Black-Scholes Using Girsanov We may assume a riskfree rate r_t and the stock price S_t satisfies:

$$dS_t = \mu_t S_t dt + \sigma_t S_t dW_t,$$

where μ_t and σ_t are adapted to \mathcal{F}_t . Using Ito's formula, we have

$$d\left(\frac{S_t}{M_t}\right) = \sigma_t \frac{S_t}{M_t} \left(\frac{\mu_t - r_t}{\sigma_t} dt + dW_t\right).$$

If we define

$$\eta_t = \frac{\mu_t - r_t}{\sigma_t}$$
$$L_t = -\int_0^t \eta_s dW_s,$$

and similarly

$$\xi_t = \exp\left(L_t - \frac{1}{2}[L]_t\right)$$

and $\tilde{\mathbb{P}}$ such that $d\tilde{\mathbb{P}} = \xi d\mathbb{P}$.

Then we have

$$d\left(\frac{S_t}{M_t}\right) = \sigma_t \frac{S_t}{M_t} d\tilde{W}_t.$$

Or equivalently,

$$dS_t = r_t S_t dt + \sigma_t S_t d\tilde{W}_t.$$

Then the price of an European call option would be

$$C_t = \tilde{\mathbb{E}}\left\{\exp\left(-\int_t^T r_s ds\right) \max(S_T - K, 0) |\mathcal{F}_t\right\} = M_t \tilde{\mathbb{E}}\left\{\frac{C_T}{M_T} |\mathcal{F}_t\right\},$$

where $\tilde{\mathbb{E}}$ is taken with respect to $\tilde{\mathbb{P}}$.

Note that the process η_t satisfies

$$\mu_t - r_t = \eta_t \sigma_t.$$

 η_t measures the excess return per unit of risk the market offers. For this reason η_t is called the "market price of risk" process.

Example: Forwards and Futures A forward contract is an agreement to pay a specified delivery price K for an asset at a delivery date T. Suppose the asset price process is S_t . At time T, the value of the contract is $S_T - K$. At the time of reaching an agreement, say t, the value of the contract must be zero,

$$M_t \tilde{\mathbb{E}}_t \left(\frac{S_T - K}{M_T} \right) = \tilde{\mathbb{E}}_t \exp\left(-\int_t^T r_s ds \right) (S_T - K) = S_t - P(t, T)K = 0,$$

where $P(t,T) = \tilde{\mathbb{E}}_t \left(\exp\left(-\int_t^T r_s ds\right) \right)$. Thus we obtain the forward price,

$$K = S_t / P(t, T).$$

Note that the forward price is a function of both t and T.

After a forward contract is signed on t, the value of this agreement will most likely diverge from zero, often substantially. Let u be such that t < u < T. For the party with the long position, who receives S_T and pays $S_t/P(t,T)$ at time T, the value of the agreement at time u is

$$M_u \tilde{\mathbb{E}}_u \left(M_T^{-1} \left(S_T - \frac{S_t}{P(t,T)} \right) \right) = S_u - S_t \frac{P(u,T)}{P(t,T)}.$$

If the riskfree rate is a constant r, then it becomes $S_u - \exp(r(u-t))S_t$.

If the asset price rises more rapidly than the money account, then the long (short) position has a positive (negative) value. If the growth rate is less than the riskfree rate, then the long (short) position has a negative (positive) value. Whichever happens, one party will have an incentive to default.

A futures contract alleviates the risk of default by margin requirement (initial margin, marking to margin) and by trading in an exchange market. Suppose the spot price process of an underlying asset is S_t and let F(t,T) denote the futures price process of a contract that matures at T. We have $F(T,T) = S_T$.

F(t,T) must be a martingale process under the risk-neutral measure \mathbb{P} . To see this, consider a partition of the life span of a futures contract, $t = t_0 < t_1 < \cdots <$

 $t_n = T$. Each interval $[t_k, t_{k+1})$ represents a "day". A long position in the futures contract is an agreement to receive as a cash flow the changes in the futures price, $F(t_{k+1}, T) - F(t_k, T)$, during the time the position is held. A short position receives the opposite. Suppose that the riskfree rate is constant within each day and that it is determined in the previous day. Then $M_{t_{k+1}}$ is \mathcal{F}_{t_k} -measurable, since

$$M_{t_{k+1}} = \exp\left(-\int_0^{t_{k+1}} r_s ds\right) = \exp\left(-\sum_{k=0}^k r_{t_k}(t_{k+1} - t_k)\right).$$

In equilibrium, any future cash flow must have a zero current value. That is, for all k,

$$M_{t_k} \tilde{\mathbb{E}}\left(M_{t_{k+1}}^{-1}\left(F(t_{k+1},T) - F(t_k,T)\right) | \mathcal{F}_{t_k}\right) = 0.$$

Since $M_{t_{k+1}}$ is \mathcal{F}_{t_k} -measurable, we have

$$\frac{M_{t_k}}{M_{t_{k+1}}} \tilde{\mathbb{E}} \left(F(t_{k+1}, T) - F(t_k, T) | \mathcal{F}_{t_k} \right) = 0.$$

Hence $F(t_k, T)$ must be a martingale sequence. And obviously, $F(t_{n-1}, T) = \mathbb{E}_{t_{n-1}}S_T$, $F(t_{n-2}, T) = \mathbb{E}_{t_{n-2}}F(t_{n-1}, T)$, and so on. By the law of iterative expectation, we have

$$F(t,T) = \mathbb{E}(S_T | \mathcal{F}_t).$$

Forwards-Futures Spread The difference between forward and futures prices,

$$D_t = S_t / P(t, T) - \mathbb{E}(S_T | \mathcal{F}_t),$$

is called the forward-futures spread. It is obvious that $D_t \to 0$ as $t \to T$.

Since
$$S_t = \tilde{\mathbb{E}}_t \left(\frac{M_t}{M_T} S_T \right)$$
 and $P(t,T) = \tilde{\mathbb{E}}_t \frac{M_t}{M_T}$, we have
 $D_t = \frac{1}{P(t,T)} \left(\tilde{\mathbb{E}}_t \left(\frac{M_t}{M_T} S_T \right) - \tilde{\mathbb{E}}_t \frac{M_t}{M_T} \tilde{\mathbb{E}}_t S_T \right) = \frac{1}{P(t,T)} \widetilde{\text{cov}} \left(\frac{M_t}{M_T}, S_T | \mathcal{F}_t \right).$

If r_t is a constant, $D_t = 0$. If P(t,T) is positively correlated with S_T , which means that a higher S_T goes together with a lower interest rate, then the forward price is higher than the futures price.

Example: Pricing of Cash Flow Suppose an asset pays D_t between time 0 and t. Then a long position of the asset gives us a gain process that satisfies

$$dG_t = dD_t + r_t G_t d_t,$$

or

$$d(G_t/M_t) = 1/M_t dD_t.$$

The risk-neutral price at time t of the cash flow between t and T is thus,

$$M_t \tilde{\mathbb{E}}_t(G_T/M_T) = M_t \tilde{\mathbb{E}}_t\left(\int_t^T 1/M_s dD_s\right).$$

The cash flow may be negative, in which case D_t is decreasing. The cash flow is most likely discrete, ie,

$$D_t = \sum_{i=1}^n d_i I_{[0,t]}(t_i),$$

where $0 < t_1 < t_2 < \cdots < t_n \leq T$ and d_i is random payment at time t_i . In this case, the risk neutral price at time t is given by

$$\sum_{i=1}^{n} I_{[t,T]}(t_i) \left(M_t \tilde{\mathbb{E}}_t(M_{t_i} d_i) \right).$$

If d_i is deterministic, then the above formula reduces to the pricing formula for bond with fixed coupons.

The Multivariate Case

We first state the multivariate Girsanov theorem. It follows easily from the general Girsanov theorem.

Multivariate Girsanov Let $\eta_t \in \mathbb{R}^d$ be adapted to \mathcal{F}_t and let

$$\xi_t = \exp\left(-\int_0^t \eta_s \cdot dW_s - \frac{1}{2}\int_0^t \|\eta_s\|^2 ds\right).$$
(3.15)

If W_t is P-BM, then $\tilde{W}_t = W_t + \int_0^t \eta_s ds$ is BM under $\tilde{\mathbb{P}}$.

We consider a stock market of N stocks. Let $S_t = (S_t^1, ..., S_t^N)$ be the stock prices and let $W_t = (W_t^1, ..., W_t^d)$ be a *d*-dimensional independent Brownian Motions.

Assume that for each stock,

$$dS_t^i = \mu_t^i S_t^i dt + S_t^i \sum_{j=1}^d \sigma_t^{ij} dW_t^j,$$

= $\mu_t^i S_t^i dt + S_t^i \sigma_t^i \cdot dW_t, \quad i = 1, ..., N,$

where $\sigma_t^i = (\sigma_t^{i1}, ..., \sigma_t^{Nd}).$

If we find an adapted process (η_t) such that

$$\mu_t^i - r_t = \eta_t \cdot \sigma_t^i, \quad i = 1, ..., N,$$
(3.16)

we may define ξ as in (3.15) and define accordingly $\tilde{\mathbb{P}}$ such that $\tilde{W}_t = W_t + \int_0^t \eta_s ds$ is BM under $\tilde{\mathbb{P}}$. Hence

$$dS_t^i = r_t S_t^i dt + S_t^i \sigma_t^i \cdot d\tilde{W}_t, \quad i = 1, ..., N.$$

Or

$$d\left(\frac{S_t^i}{M_t}\right) = \left(\frac{S_t^i}{M_t}\right)\sigma_t^i \cdot d\tilde{W}_t.$$

In other words, each numeraire-denominated stock price is a martingale under \mathbb{P} .

We restate the crucial condition in (3.16) in matrix form. Let $\mu_t = (\mu_t^1, ..., \mu_t^N)'$, we have

$$\sigma_t \eta_t = \mu_t - r_t. \tag{3.17}$$

Recall that (η_t) is called the "market-price-of-risk" (MPR) process and measures the drift in price the investor get compensated for taking each unit of risk.

When the equation (3.17) has no solution, martingale equivalent measure for this market does not exist. In this case, it is always possible to find arbitrage strategies. This says that no arbitrage implies the existence of the "market-priceof-risk" process, hence the existence of martingale equivalent measure.

When N > d, some of the securities are "redundant" (derivatives, for example) and can be replicated by a linear combination of other stocks. Thus we may assume N = d. If rank $(\sigma) = d$, there is at most one MPR process, and accordingly, an equivalent martingale measure. The market is said to be complete.

Market Completeness Let $\mathcal{M} = \{h_T \cdot X_T | (h_t) \text{ is self-financing} \}$. If $\mathcal{M} = \mathcal{L}^2$, the space of all finite-variance random variables, we say the market is *complete*.

A necessary and sufficient condition for complete market is that there is an MPR process and rank(σ_t) = d. In other words, there exists a unique MPR process.

Hedging a General Contingent Claim

We first state an important theorem.

Theorem 3.4.1 (Martingale Representation Theorem) Let $W = (W^1, W^2, ..., W^d)$, and \mathcal{F}_t be the natural filtration of W. If (M_t) is a martingale w.r.t. \mathcal{F}_t , then there exists $K = (K^1, K^2, ..., K^d)$ such that $\int_0^t (K_s^j)^2 ds < \infty$ for each j and

$$M_t = M_0 + \int_0^t K_s \cdot dW_s.$$

Given a \mathcal{F}_T -measurable contingent claim C_T , the no arbitrage price at time t satisfies,

$$\frac{C_t}{M_t} = \tilde{\mathbb{E}}\left(\frac{C_T}{M_T} | \mathcal{F}_t\right).$$

The process of $(C_t/M_t, \mathcal{F}_t)$ is a $\tilde{\mathbb{P}}$ martingale. By martingale representation theorem, we can find an adapted process (γ_t) such that

$$\frac{C_t}{M_t} = C(0) + \int_0^t \gamma_s d\tilde{W}_s.$$

Let the value of the hedging portfolio be H_t . Suppose we hold Δ_t amount of stock at time t, we have

$$dH_t = \Delta_t dS_t + r_t (H_t - \Delta_t S_t) dt$$

= $r_t H_t dt + \Delta_t \sigma_t S_t (\eta_t dt + dW_t),$

where η_t is the market price of risk process. Hence

$$d\left(\frac{H_t}{M_t}\right) = \Delta_t \sigma_t S_t(\eta_t dt + dW_t) = \Delta_t \sigma_t S_t M_t^{-1} d\tilde{W}_t.$$

So we have

$$\left(\frac{H_t}{M_t}\right) = H_0 + \int_0^t \Delta_v \sigma_v S_v M_v^{-1} d\tilde{W}_v.$$

To hedge C_t , we must have $H_0 = C_0$ and

$$\Delta_t = \frac{M_t}{\sigma_t S_t} \gamma_t.$$

From Risk Neutral Pricing to PDE

Consider the Black-Scholes Model,

$$dM_t = r_t M_t dt$$

$$dS_t = \mu_t S_t dt + \sigma_t S_t dW_t$$

The MPR η exists,

$$\eta_t = \frac{\mu_t - r_t}{\sigma_t}.$$

And σ is trivially full rank. Hence the market is complete.

Since $S_T \in \mathcal{L}^2$, $C_T = \max(S_T - K, 0) \in \mathcal{L}^2$. We know that $M_t^{-1}C_t = \tilde{\mathbb{E}}_t M_T^{-1}C_T$. Let $F(S_t, t) = C_t$, we have

$$d\left(\frac{F(S_t,t)}{M_t}\right) = \frac{1}{M_t}(-r_t F(S_t,t)dt + dF(S_t,t))$$

= $\frac{1}{M_t}(F_2 + r_t S_t F_1 + \frac{1}{2}F_{11}\sigma_t^2 S_t^2 - r_t F)dt + \frac{S_t}{M_t}\sigma_t F_1 d\tilde{W}_t.$

The bounded variation part gives the PDE. And the martingale part gives the hedging strategy, which is

$$\Delta_t = F_1(S_t, t).$$

A complete market admits only one martingale equivalent measure, in which case C_t is unique.

3.5 State Prices

Definition

A state-price deflator is a deflator m for a price process X such that X^m is a martingale w.r.t. the natural filtration.

Other names for m are stochastic discount factor, state-price density, marginal rates of substitution, and pricing kernel.

Given a numeraire M_t and an equivalent martingale measure ξ , the state-price deflator is

$$m_t = \frac{\xi_t}{M_t}$$

To see this,

$$\mathbb{E}_s m_t X_t = \mathbb{E}_s \xi_t \frac{X_t}{M_t} = \xi_s \tilde{\mathbb{E}}_s \frac{X_t}{M_t} = \xi_s \frac{X_s}{M_s} = m_s X_s$$

Conversely, given m_t , we can construct ξ_t by

$$\xi_t = M_t m_t.$$

Risk Premium

 m_t is an Ito process, hence it can be characterized by

$$dm_t = m_t \mu_{m,t} dt + m_t (\sigma_{m,t} \cdot dW_t).$$

Since

$$dm_t = d\left(\frac{\xi_t}{M_t}\right) = -r_t m_t dt + M_t^{-1} d\xi_t,$$

 So

$$\mu_{m,t} = -r_t.$$

Let a price process be S_t^i ,

$$dS_t^i = S_t^i \mu_t^i dt + S_t^i (\sigma_t^i \cdot dW_t).$$

We have

$$\begin{aligned} d(m_t S_t^i) &= m_t dS_t^i + S_t^i dm_t + d[m, S^i]_t \\ &= m_t S_t^i (\mu_t^i + \mu_{m,t} + \sigma_{m,t} \cdot \sigma_t^i) dt + m_t S_t^i (\sigma_{m,t} + \sigma_t^i) \cdot dW_t. \end{aligned}$$

 $(m_t S_t)$ is a martingale, so

$$\mu_t^i + \mu_{m,t} + \sigma_{m,t} \cdot \sigma_t^i = 0.$$

$$\mu_t^i - \pi - \sigma_t^i - \sigma_t^i$$

Hence,

$$\mu_t^i - r_t = -\sigma_t^i \cdot \sigma_{m,t}, \tag{3.18}$$

where both σ_t and $\sigma_{m,t}$ can be negative. (3.18) characterizes the excess expected return or risk premium of the stock S^i .

Furthermore, if we define

$$\beta_t^i = -\frac{\sigma_t^i \cdot \sigma_{m,t}}{\sigma_{m,t} \cdot \sigma_{m,t}} \text{ and } \lambda_{m,t} = \sigma_{m,t} \cdot \sigma_{m,t} \equiv \|\sigma_{m,t}\|^2,$$

then we have

$$\mu_t^i = r_t + \beta_t^i \lambda_{m,t}$$

 β_t^i measures the systematic risk in S_t^i , and $\lambda_{m,t}$ measures the price of the systematic risk. Note that $\eta_t = -\sigma_{m,t}$, since

$$\eta_t \cdot \sigma_t^i = \mu_t^i - r_t = -\sigma_{m,t} \cdot \sigma_t^i$$
 for all t .

So

$$\lambda_{m,t} = \|\eta_t\|^2.$$

3.6 Treatment of Dividends

We discuss how to treat dividend payment in risk-neutral pricing framework.

Continuous Payment

We assume that if a stock withholds dividends, the stock price follows a diffusion process,

$$dS_t = \mu_t S_t dt + \sigma_t S_t dW_t.$$

Now, we may assume that the stock pays dividends continuously at a rate of d_t . Then

$$dS_t = \mu_t S_t dt + \sigma_t S_t dW_t - d_t S_t dt.$$

To replicate the price process of a contingent claim, $(C_t, t \in [0, T])$, we construct a self-financing portfolio h_t which holds Δ_t stocks at time t. Let H_t be the value of the portfolio and let D_t denote the dividends paid cumulatively up to time t and satisfies

$$dD_t = d_t S_t dt$$

Then we have

$$dH_t = \Delta_t dS_t + \Delta_t dD_t + r_t (H_t - \Delta_t S_t) dt$$

= $r_t H_t dt + \Delta_t S_t (\mu_t - r_t) dt + \Delta_t S_t \sigma_t dW_t$
= $r_t H_t dt + \sigma_t \Delta_t S_t (\eta_t dt + dW_t),$

where

$$\eta_t = \frac{\mu_t - r_t}{\sigma_t}$$

is the MPR process. Define $d\tilde{W}_t = dW_t + \eta_t dt$. Under $\tilde{\mathbb{P}}$, density process of which is defined as the exponential martingale of $L_t = -\int_0^t \eta_s dW_s$, we have

$$dH_t = r_t H_t dt + \sigma_t \Delta_t S_t dW_t.$$

In other words, under $\tilde{\mathbb{P}}$, the numeraire-deflated process H_t/M_t is a martingale,

$$d\left(\frac{H_t}{M_t}\right) = \sigma_t \Delta_t S_t / M_t d\tilde{W}_t$$

Hence the price of the contingent claim would be given by

$$C_t = M_t \tilde{\mathbb{E}}_t \frac{C_T}{M_T}.$$

It can be shown that under $\tilde{\mathbb{P}}$, the stock price follows

$$dS_t = (r_t - d_t)S_t dt + \sigma_t S_t dW_t.$$

Now let $d_t = d$, $r_t = r$, and $\sigma_t = \sigma$, we have

$$S_T = S_0 \exp\left[(r - d - \frac{1}{2}\sigma^2)T + \sigma \tilde{W}_T\right].$$

From this we can easily calculate Black-Scholes Formula with continuous dividend yield d.

A Different Perspective

Obviously, if we reinvest the dividends, the "gain process" of the stock follows

$$dG_t = \mu_t G_t dt + \sigma_t G_t dW_t.$$

Consider the market that consists of the stock with dividend reinvestment and a money account, $X_t = (G_t, M_t)$. This market admits no arbitrage if and only if X_t/M_t admits an equivalent martingale measure, say, $\tilde{\mathbb{P}}$. Then the price of a contingent claim $(C_t, t \in [0, T])$ in the unit of a numeraire should be a martingale under $\tilde{\mathbb{P}}$.

To find such a probability measure, we obtain

$$d\left(\frac{G_t}{M_t}\right) = (\mu_t - r_t)G_t/M_t dt + \sigma_t G_t/M_t dW_t = \sigma_t G_t/M_t \left(\frac{\mu_t - r_t}{\sigma_t} + dW_t\right).$$

Define $\tilde{W}_t = W_t + \int_0^t \eta_s ds$, $\eta_t = (\mu_t - r_t)/\sigma_t$, $L_t = -\int_0^t \eta_s dW_s$, and $\xi_t = \exp(L_t - [L]_t/2)$, which corresponds to an equivalent probability measure $\tilde{\mathbb{P}}$. According to Girsanov theorem, $\tilde{W}_t \sim \tilde{\mathbb{P}}$ -Brownian Motion. Then G_t/M_t is $\tilde{\mathbb{P}}$ -martingale.

Discrete Payment

Suppose the dividends are paid at n time points on [0, T], $0 < t_1 < t_2 < ... < t_n < T$. At each time point t_i , the dividend payment is $d_i S(t_{i-})$, where d_i is \mathcal{F}_{t_i} -measurable and $S(t_{i-})$ denotes the stock price just prior to the payment. The stock price after the payment is

$$S_{t_i} = S_{t_i-} - d_i S_{t_i-} = (1 - d_i) S_{t_i-}.$$

We assume that between dividend payment dates the stock price follows,

$$dS_t = \mu_t S_t dt + \sigma_t S_t dW_t, \quad t_i \le t < t_{i+1}.$$

Hence, for time $t \in [t_i, t_{i+1})$, the value of hedging portfolio H_t follows

$$dH_t = \Delta_t dS_t + (H_t - \Delta_t S_t) dM_t$$

= $r_t H_t dt + \Delta_t S_t (\mu_t - r_t) dt + \Delta_t S_t \sigma_t dW_t$
= $r_t H_t dt + \sigma_t \Delta_t S_t (\eta_t dt + dW_t),$

where

$$\eta_t = \frac{\mu_t - r_t}{\sigma_t}$$

Since the portfolio collects the dividend payment, the portfolio value does not jump at payment dates. Hence the above SDE describes the portfolio value for all t.

We may define $d\tilde{W}_t = dW_t + \eta_t dt$ and define $\tilde{\mathbb{P}}$ as usual. Then H_t/M_t would be martingale under $\tilde{\mathbb{P}}$ for all t. And

$$dS_t = r_t S_t dt + \sigma_t S_t dW_t, \ t \in [t_i, t_{i+1}), \ i = 0, ..., n$$

Now let $r_t = r$ and $\sigma_t = \sigma$, we have

$$S_{t_{i+1}-} = S_i \exp\left[(r - \frac{1}{2}\sigma^2)(t_{i+1} - t_i) + \sigma(\tilde{W}_{t_{i+1}} - \tilde{W}_{t_i})\right],$$

and

$$S_{t_{i+1}} = (1 - d_{i+1})S_i \exp\left[(r - \frac{1}{2}\sigma^2)(t_{i+1} - t_i) + \sigma(\tilde{W}_{t_{i+1}} - \tilde{W}_{t_i})\right].$$

Then we have

$$S_T = \left(S_0 \prod_{i=0}^{n-1} (1 - d_{i+1})\right) \cdot \exp\left[(r - \frac{1}{2}\sigma^2)T + \sigma \tilde{W}_T\right].$$

It is easy to see that we can use the Black-Scholes formula to calculate the price of European call options on the stock S_t , with initial value being replaced by $(S_0 \prod_{i=0}^{n-1} (1 - d_{i+1}))$.

Chapter 4

Term Structure Modeling

4.1 Basics

We study the price of money, i.e., the default-free interest rate. (Think of the interest rate on bills/notes/bonds issued by the US treasury.) This differs from the price of a particular bond in that the latter depends on factors other than the time value of money, such as the credit history of the borrower, the liquidity condition of the market, and so on.

Term Structure

We assume that a continuum of default-free discount bonds trade continuously at time t with differing maturities T and prices P(t,T). Assume P(T,T) = 1. P(t,T) is called the term structure.

P(t,T) can be read along two dimensions:

- 1. Fix t and let T vary: prices for different maturities at time t.
- 2. Fix T and let t vary: historical price series of a particular maturity T.

Yield Curve

The interest rate implied by the zero-coupon bond is called spot rate, which is given by

$$R(t,T) = -\frac{\log P(t,T)}{T-t}.$$

Fixing a t and varying T, we call R(t,T) the yield curve. Yield curves can be increasing or decreasing functions of T. In practice, yields to maturity on coupon bonds are often calculated. The yield to maturity is the internal rate of return for a coupon bond, or, the constant interest rate that makes the present value of future cash flow (coupon payments and the face value) equal to the market price of the bond. Suppose the coupon bond in question pays a series of coupons in the remaining period (c_i at $t < t_i \leq T$, including the principal), then the yield to maturity (YTM), y(r, T), solves the following equation,

$$P_c(t,T) = \sum_{t_i > t} c_i \exp(-(t_i - t)y(t,T)), \qquad (4.1)$$

where $P_c(t, T)$ is the market price of the coupon bond. Obviously YTM is not only a function of T, but also how coupons are paid (annual, biannual, or quarterly). For coupon bonds, the duration of a coupon bond is defined as:

$$D_t = \frac{\sum_{t_i > t} (t_i - t) c_i \exp(-(t_i - t)y(t, T))}{\sum_{t_i > t} c_i \exp(-(t_i - t)y(t, T))}.$$

The duration D_t thus defined is in fact the derivative of $-\log P_c(t,T)$ with respect to y(t,T). Intuitively, that is the percentage decline of the bond price associated with a unit increase in YTM. For example, the price of a 5-year-duration bond declines 5% when its YTM increases by 1%. For discount bonds, duration is exactly the term to maturity (TTM), that is (T-t). Duration may be understood as a measure of risk for coupon bonds.

Short Rate

The short rate, or the instantaneous rate, measures the *current* cost of short-term borrowing.

$$r_t = \lim_{\Delta \to 0} R(t, t + \Delta) = -\lim_{\Delta \to 0} \frac{\log(P(t, t + \Delta))}{\Delta}.$$

Or,

$$r_t = R(t,t)$$
, and
 $r_t = -\frac{\partial}{\partial T} \log P(t,t)$

Forward Rate

Let $t < T_1 < T_2$. Consider a forward contract on a bond that matures at T_2 : an agreement at time t to make a payment at T_1 and receive a payment in return at T_2 .
We can replicate the contract, at time t, by

- buying a T_2 bond
- selling k units of T_1 bond.

The cash flow of this portfolio is

- At $t, -P(t, T_2) + kP(t, T_1)$
- At T_1 , -k
- At T_2 , 1

Since the value of any forward contract should be zero at the time of agreement t, k must satisfies

$$k = \frac{P(t, T_2)}{P(t, T_1)}.$$

Obviously, k should be called the forward price of the T_2 -bond. The corresponding yield of holding the T_2 -bond in the interval of $[T_1, T_2]$, denoted as $F(t, T_1, T_2)$, is

$$F(t, T_1, T_2) = \frac{\log 1/k}{T_2 - T_1} = -\frac{\log P(t, T_2) - \log P(t, T_1)}{T_2 - T_1}.$$

Now it is ready to define forward rate, the forward price for instantaneous borrowing at time T,

$$f(t,T) = \lim_{T_2 \to T} F(t,T,T_2) = \lim_{\Delta \to 0} -\frac{\log P(t,T+\Delta) - \log P(t,T)}{\Delta} = -\frac{\partial}{\partial T} \log P(t,T).$$

The forward rate f(t,T) contains all information about P(t,T) and R(t,T). Specifically, we have

$$P(t,T) = \exp\left(-\int_{t}^{T} f(t,u)du\right),$$

and

$$R(t,T) = \frac{\int_{t}^{T} f(t,u)du}{T-t}$$

And the short rate r_t can be recovered using

$$r_t = f(t, t).$$

We also have

$$\frac{\partial R(t,T)}{\partial T} = -\frac{\partial \log P(t,T)}{\partial T} \frac{1}{T-t} + \frac{\log P(t,T)}{(T-t)^2}$$
$$= \frac{f(t,T)}{T-t} - \frac{1}{T-t}R(t,T).$$

Hence

$$f(t,T) = R(t,T) + (T-t)\frac{\partial R(t,T)}{\partial T}.$$

When T = t, $f(t,t) = R(t,t) = r_t$. Otherwise, f(t,T) is greater (less) than R(t,T) when R(t,T) is increasing (decreasing). Finally, we have

$$F(t, T_1, T_2) = \frac{1}{T_2 - T_1} \int_{T_1}^{T_2} f(t, s) ds.$$

4.2 The Single-Factor Heath-Jarrow-Morton Model

The Risk-Neutral Pricing

The Model P(t,T), R(t,T), and f(t,T) contain the same information. The HJM model (Heath, Jarrow, and Morton, 1992) is a model on f(t,T):

$$f(t,T) = f(0,T) + \int_0^t \alpha(s,T) ds + \int_0^t \sigma(s,T) dW_s.$$
 (4.2)

Or, in its differential form,

$$d_t f(t,T) = \alpha(t,T)dt + \sigma(t,T)dW_t.$$

- $\alpha(t,T)$ and $\sigma(t,T)$ may depend on $(W_s, s \leq t)$ and f(t,T) itself.
- f(0,T) is deterministic.
- $\int_0^T \int_0^u |\alpha(t,u)| dt du < \infty$ and $\mathbb{E}\left(\int_0^T \left|\int_0^u \sigma(t,u) dW_t\right| du\right) < \infty$.

The Numeraire We use M_t , which satisfies

$$M_0 = 1$$
, and $dM_t = r_t M_t dt$.

Or

$$M_t = \exp\left(\int_0^t r_s ds\right).$$

Recall that $r_t = f(t, t)$. So

$$r_t = f(0,t) + \int_0^t \alpha(s,t)ds + \int_0^t \sigma(s,t)dW_s.$$

 So

$$M_t = \exp\left(\int_0^t f(0,s)ds + \int_0^t \int_0^u \alpha(s,u)dsdu + \int_0^t \int_0^u \sigma(s,u)dW_sdu\right)$$
$$= \exp\left(\int_0^t f(0,s)ds + \int_0^t \int_s^t \alpha(s,u)duds + \int_0^t \int_s^t \sigma(s,u)dudW_s\right).$$

The Bond We can choose any bond price to construct martingale equivalent measure under no arbitrage condition. Consider P(t,T),

$$P(t,T) = \exp\left(-\int_{t}^{T} f(t,u)du\right)$$

= $\exp\left(-\left[\int_{t}^{T} f(0,u)du + \int_{0}^{t} \int_{t}^{T} \alpha(s,u)duds + \int_{0}^{t} \int_{t}^{T} \sigma(s,u)dudW_{s}\right]\right)$

We can check that $P(0,T) = \exp\left(-\int_0^T f(0,u)du\right)$, and P(T,T) = 1.

Deflation Define

$$Z(t,T) = M_t^{-1}P(t,T)$$

= $\exp\left[-\int_0^T f(0,u)du - \int_0^t \int_s^T \alpha(s,u)duds - \int_0^t \left(\int_s^T \sigma(s,u)du\right)dW_s\right]$
= $\exp\left[-\int_0^T f(0,u)du - \int_0^t \int_s^T \alpha(s,u)duds + \int_0^t \Sigma(s,T)dW_s\right],$

where $\Sigma(t,T) = -\int_t^T \sigma(t,u) du$. Let X_t be the term in the bracket,

$$d_t Z(t,T) = d(\exp(X_t))$$

= $\exp(X_t) dX_t + \frac{1}{2} \exp(X_t) d[X]_t$
= $Z(t,T) \left(\left(\frac{1}{2} \Sigma(t,T)^2 - \int_t^T \alpha(t,u) du \right) dt + \Sigma(t,T) dW_t \right).$

Change of Measure Then the market-price-of-risk process would be

$$\eta_t = \frac{\frac{1}{2}\Sigma^2(t,T) - \int_t^T \alpha(t,u)du}{\Sigma(t,T)}$$
$$= \frac{1}{2}\Sigma(t,T) - \Sigma^{-1}\int_t^T \alpha(t,u)du.$$

Then we can define $\tilde{\mathbb{P}}$ and a \tilde{W}_t such that

$$dW_t = dW_t + \eta_t dt.$$

Then

$$d_t Z(t,T) = Z(t,T)\Sigma(t,T)d\tilde{W}_t.$$
(4.3)

Hence Z(t,T) is a $\tilde{\mathbb{P}}$ -martingale. And under $\tilde{\mathbb{P}}$, the bond price P(t,T) has a drift term r_t :

$$d_t P(t,T) = P(t,T)(r_t dt + \Sigma(t,T) dW_t).$$

Other Bonds We use P(t,T) to construct a martingale equivalent measure \mathbb{P} . What about other bonds, such as P(t,S), S < T?

Let X = 1 be a claim that pays off at time S. Then P(t, S) is the price of X at time t,

$$P(t,S) = M_t \tilde{\mathbb{E}}_t(M_S^{-1}) = \tilde{\mathbb{E}}_t \left(\exp\left(-\int_t^S r_s ds\right) \right).$$

And the deflated price process is

$$Z(t,S) = M_t^{-1} P(t,S) = \tilde{\mathbb{E}}_t(M_S^{-1}).$$

So the deflated prices of all other bonds are $\tilde{\mathbb{P}}$ -martingale. This means that their $\tilde{\mathbb{P}}$ -drifts are restricted such that η_t is the same market-price-of-risk process for all bonds. In particular, for all $S \in [0, T]$,

$$\int_{t}^{S} \alpha(t,s) ds = \frac{1}{2} \Sigma^{2}(t,S) - \Sigma(t,S) \eta_{t}.$$

Taking $\partial/\partial S$ on both sides,

$$\alpha(t,S) = -\Sigma(t,S)\sigma(t,S) + \sigma(t,S)\eta_t$$

= $\sigma(t,S)(\eta_t - \Sigma(t,S)).$

If there exists an (η_t) such that the above holds, among other regularity conditions, then the market is complete. We may find a self-financing portfolio of $(M_t, P(t, T))$, that replicates any contingent claim U_S which pays off at time S < T.

A Direct Approach

We have

$$f(t,T) = f(0,T) + \int_0^t \alpha(s,T)ds + \int_0^t \sigma(s,T)(d\tilde{W}_s - \eta_s ds)$$

= $f(0,T) + \int_0^t (\alpha(s,T) - \sigma(s,T)\eta_s)dt + \int_0^t \sigma(s,T)d\tilde{W}_s$
= $f(0,T) + \int_0^t (-\Sigma(s,T)\sigma(s,T))dt + \int_0^t \sigma(s,T)d\tilde{W}_s.$

So as in Duffie (2001), we may directly assume no arbitrage opportunities and there exists a martingale equivalent measure $\tilde{\mathbb{P}}$ such that we may specify f(t,T) as

$$f(t,T) = f(0,T) + \int_0^t \mu(s,T) ds + \int_0^t \sigma(s,T) d\tilde{W}_s,$$

where

$$\mu(t,T) = -\sigma(t,T)\Sigma(t,T)$$
$$= \sigma(t,T) \int_t^T \sigma(t,s) ds$$

Then

$$r_t = f(0,t) + \int_0^t \sigma(s,t) \int_s^t \sigma(s,u) du ds + \int_0^t \sigma(s,T) d\tilde{W}_s.$$

4.3 Short-Rate Models

The short-rate model is a model on r_t . Assume there exists a martingale equivalent measure $\tilde{\mathbb{P}}$. r_t is usually specified as a Markov diffusion:

$$dr_t = \nu(r_t)dt + \rho(r_t)d\tilde{W}_t.$$
(4.4)

Then the term structure P(t,T) is given as

$$P(t,T) = \tilde{\mathbb{E}}_t \left(\exp(-\int_t^T r_s ds) \right).$$

Note that the short rate process alone does not recover the term structure, which is determined by the risk appetite of the market as well as the future short-term borrowing cost.

Connection with Forward-Rate Models

Given f(t, T), we can easily recover r_t . And we may recover the forward rate f(t, T) from the short rate r_t as follows. We define g(x, t, T) as

$$g(x,t,T) = -\log\left[\tilde{\mathbb{E}}\left(\exp\left(-\int_{t}^{T} r_{s} ds\right) \mid r_{t} = x\right)\right]$$

Then we have

$$g(r_t, t, T) = -\log P(t, T) = \int_t^T f(t, u) du.$$

In other words,

$$f(t,T) = \frac{\partial}{\partial T}g(r_t, t, T)$$

Hence,

$$d_t f(t,T) = \frac{\partial^2 g}{\partial T \partial t} dt + \frac{\partial^2 g}{\partial T \partial x} dr_t + \frac{1}{2} \frac{\partial^3 g}{\partial T \partial x^2} d[r]_t \\ = \left(\frac{\partial^2 g}{\partial T \partial t} \nu(r_t) + \frac{\partial^2 g}{\partial T \partial t} + \frac{1}{2} \frac{\partial^3 g}{\partial T \partial x^2} \rho^2(r_t) \right) dt + \frac{\partial^2 g}{\partial x \partial T} \rho(r_t) d\tilde{W}_t.$$

 So

$$\sigma(t,T) = \frac{\partial^2 g(r_t,t,T)}{\partial T \partial x} \rho(r_t)$$

$$\Sigma(t,T) = -\frac{\partial g(r_t,t,T)}{\partial x} \rho(r_t).$$

Note also that

$$f(0,T) = \frac{\partial g(r_0,0,T)}{\partial T}.$$

f(0,T) together with $\sigma(t,T)$ determines f(t,T) under $\tilde{\mathbb{P}}$.

Examples

Ho and Lee Model The short rate process r_t satisfies

$$dr_t = \nu_t dt + \rho d\tilde{W}_t, \tag{4.5}$$

where ν_t is deterministic and bounded and ρ is constant.

For $s \geq t$,

$$r_s = r_t + \int_t^s \nu_u du + \int_t^s \rho d\tilde{W}_u.$$

Hence

$$\int_{t}^{T} r_{s} ds = r_{t}(T-t) + \int_{t}^{T} \int_{t}^{s} \nu_{u} du ds + \int_{t}^{T} \int_{t}^{s} \rho d\tilde{W}_{u} ds$$
$$= r_{t}(T-t) + \int_{t}^{T} \int_{u}^{T} \nu_{u} ds du + \int_{t}^{T} \int_{u}^{T} \rho ds d\tilde{W}_{u}$$
$$= r_{t}(T-t) + \int_{t}^{T} \nu_{u}(T-t) du + \int_{t}^{T} \rho(T-u) d\tilde{W}_{u}$$

Let $M_T = -\int_t^T \rho(T-u) d\tilde{W}_u$. M_T is a $\tilde{\mathbb{P}}$ -martingale and $M_t = 0$. We have

$$[M]_T = \rho^2 \int_t^T (T-u)^2 du.$$

Then $\exp(M_T - \frac{1}{2}[M]_T)$ is a (positive) $\tilde{\mathbb{P}}$ -martingale with $\exp(M_t - \frac{1}{2}[M]_t) = 1$. Hence

$$\tilde{\mathbb{E}}_t \exp(M_T) = \exp\left(\frac{1}{2}[M]_T\right) \tilde{\mathbb{E}}_t \exp\left(M_T - \frac{1}{2}[M]_T\right) \\ = \exp\left(\frac{1}{2}[M]_T\right) \exp(M_t - \frac{1}{2}[M]_t) \\ = \exp\left(\frac{1}{2}\rho^2 \int_t^T (T-u)^2 du\right).$$

We have

$$\exp\left(-\int_{t}^{T} r_{s} ds\right) = \exp\left(-r_{t}(T-t)\right) \exp\left(-\int_{t}^{T} \nu_{u}(T-u) du\right) \exp\left(M_{T}\right)$$

Hence

$$\tilde{\mathbb{E}}\left[e^{-\int_t^T r_s ds} | r_t = x\right] = e^{-x(T-t)} \cdot e^{-\int_t^T \nu_u (T-u) du} \cdot e^{\frac{1}{2}\rho^2 \int_t^T (T-u)^2 du}.$$

Hence

$$g(x,t,T) = -\log\left(\tilde{\mathbb{E}}\left(e^{-\int_t^T r_s ds} | r_t = x\right)\right)$$

= $x(T-t) + \int_t^T \nu_u(T-u) du - \frac{1}{2}\rho^2 \int_t^T (T-u)^2 du$
= $x(T-t) + \int_t^T \nu_u(T-u) du - \frac{1}{6}\rho^2 (T-t)^3.$

The HJM volatility $\sigma(t,T)$ is then

$$\sigma(t,T) = \rho \frac{\partial^2 g(r_t, t, T)}{\partial x \partial T} = \rho,$$

which does not depend on t or T.

And $\Sigma(t,T)$ is

$$\Sigma(t,T) = -\rho(T-t).$$

So Ho and Lee model is equivalent to the following HJM model:

$$d_t f(t,T) = \rho^2 (T-t)dt + \rho d\tilde{W}_t,$$

with

$$f(0,T) = \frac{\partial g(r_0, 0, T)}{\partial T} = r_0 - \frac{1}{2}\rho^2 T^2 + \int_0^T \nu_s ds$$

We may easily generalize (4.5) as follows,

$$dr_t = \nu_t dt + \rho_t dW_t.$$

The HJM counterpart would be

$$d_t f(t,T) = \rho_t^2 (T-t) dt + \rho_t d\tilde{W}_t,$$

with

$$f(0,T) = r_0 - \int_t^T \rho_s^2 (T-s) ds + \int_0^T \nu_s ds.$$

Now the HJM volatility depends on time t.

Vasicek Model Now we allow the drift depend on r_t itself,

$$dr_t = (\theta - \alpha r_t)dt + \rho d\tilde{W}_t, \tag{4.6}$$

where θ , α , and ρ are constants. The process described by (4.6) is the well-known Orstein-Uhlenbeck process. This model translates into a HJM model with volatility that depends on maturity T as well as time t.

Exercise: Show that the HJM representation of the Vasicek model is

$$\sigma(t,T) = \rho \exp(-\alpha(T-t)),$$

with

$$f(0,T) = \theta/\alpha + e^{-\alpha T}(r_0 - \theta/\alpha) - \frac{\rho^2}{2\alpha^2}(1 - e^{-\alpha T})^2.$$

We may generalize (4.6) into the following form,

$$dr_t = (\theta_t - \alpha_t r_t)dt + \rho_t dW_t,$$

where θ_t , α_t , and ρ_t are deterministic processes.

Cox-Ingersoll-Ross Model Both Ho and Lee model and Vasicek model may display negative short rates. The Cox-Ingersoll-Ross model avoids this problem.

$$dr_t = (\theta - \alpha r_t)dt + \rho \sqrt{r_t} d\tilde{W}_t, \qquad (4.7)$$

where θ , α , and ρ are constants. If $\theta \ge \rho^2/2$, r_t is positive a.s. The process described by (4.7) is the Feller's Square Root Process. We may easily generalize the CIR model to allow deterministic processes θ_t , α_t and ρ_t in place of the corresponding constants.

The HJM equivalent model needs a special function B(t,T) which is the solution to the Riccati differential equation

$$\frac{\partial B(t,T)}{\partial t} - \alpha B(t,T) - \frac{1}{2}\rho^2 B^2(t,T) + 1 = 0, \text{ with } B(T,T) = 0.$$

Then we have

$$g(x,t,T) = xB(t,T) + \theta \int_t^T B(s,T)ds.$$

Define $D(t,T) = \partial B / \partial T$. Then the HJM volatility can be written as

$$\sigma(t,T) = \rho \sqrt{r_t} D(t,T)$$

$$\Sigma(t,T) = -\rho \sqrt{r_t} B(t,T).$$

The initial value can also be easily calculated,

$$f(0,T) = r_0 D(0,T) + \theta \int_0^T D(s,T) ds.$$

Black-Karasinski Model The Black-Karasinski model forces the short rate to be positive by taking exponential of an Orstein-Uhlenbeck process:

$$r_t = \exp(X_t),\tag{4.8}$$

where

$$dX_t = (\theta_t - \alpha_t X_t)dt + \rho_t dW_t.$$

Using Ito's formula, we may write the Black-Karasinski Model as

$$dr_t = \left((\theta_t - \alpha_t \log r_t) r_t + \frac{1}{2} \rho_t^2 r_t \right) dt + \rho_t r_t d\tilde{W}_t.$$

The General Parametric Model In general, we may write the short rate model in the following form,

$$dr_t = [c_0(t) + c_1(t)r_t + c_2(t)r_t \log r_t] dt + [d_0(t) + d_1(t)r_t]^v d\tilde{W}_t$$
(4.9)

Here are some special cases.

- $c_2 = 0, d_0 = 0, v = 0.5, \text{CIR}$
- $c_1 = 0, c_2 = 0, d_1 = 0, v = 1$, Ho and Lee
- $c_2 = 0, d_1 = 0, v = 1$, Vasicek
- $d_0 = 0, v = 1$, Black-Karasinski

In particular, c_1 is usually called "mean-reversion" parameter.

If $c_0 = d_1 = 0$, r_t is Gaussian. It can be shown that g(x, t, T) satisfies

$$g(x,t,T) = A(t,T) + B(t,T)x.$$

Affine Models

When g(x, t, T) is affine in x, ie,

$$g(x, t, T) = A(t, T) + B(t, T)x,$$

then we call the associated term structure model an affine term structure model.

The term structure is affine if and only if ν and ρ^2 in the definition of short-rate models (4.4) is affine, ie,

$$\begin{aligned} \nu(r_t) &= c_0(t) + c_1(t)r_t \\ \rho^2(r_t) &= d_0(t) + d_1(t)r_t. \end{aligned}$$

Given ν and ρ , we may recover A and B. First, B satisfies the following Riccati equation,

$$\frac{\partial B(t,T)}{\partial t} + c_1(t)B(t,T) - \frac{1}{2}d_1(t)B^2(t,T) + 1 = 0, \text{ with } B(T,T) = 0.$$

And

$$A = \int_{t}^{T} \left(c_0(s)B(s,T) - \frac{1}{2}d_0(s)B^2(s,T) \right) ds.$$

In particular, Gaussian models and the CIR model have explicit solution. Others may be solved numerically.

The Feynman-Kac Formulation

In a single-factor model, the evolution of short rate depends on one factor only. To emphasize this point, we may write the short rate model in (4.4) as

$$r_t = X_t$$
, and $dX_t = \nu(X_t)dt + \rho(X_t)d\tilde{W}_t$. (4.10)

We will see that this formulation extends easily to multi-factor models.

By the Markovian nature of r_t , P(t,T) can be represented as $P(t,T) = F(r_t,t)$. Recall that

$$F(x,t) = \tilde{\mathbb{E}}_t \left[\exp\left(-\int_t^T r_s ds\right) | X_t = x \right]$$
$$= \tilde{\mathbb{E}}_t \left[\exp\left(-\int_t^T r_s ds\right) | r_t = x \right].$$

It is clear that F solves the following partial differential equation,

$$F_2(x,t) + \nu(x)F_1(x,t) + \frac{1}{2}\rho^2(x)F_{11}(x,t) - xF(x,t) = 0$$
(4.11)

with

$$F(x,T) = 1.$$

We can thus solve the above pde numerically for F(x, t), and thus P(t, T).

4.4 Multi-factor Models

Now we assume the economy is subject to more than one "shocks". Let $W = (W^1, ..., W^d)'$ be a *d*-dimensional Brownian Motion.

Multi-factor Heath-Jarrow-Morton Model

Let $\sigma = (\sigma_1, ..., \sigma_d)'$, we can specify the forward rate as

$$f(t,T) = f(0,T) + \int_0^t \alpha(s,T)ds + \int_0^t \sigma(s,T) \cdot dW_s$$

= $f(0,T) + \int_0^t \alpha(s,T)ds + \sum_{i=1}^d \int_0^t \sigma_i(s,T)dW_s^i$.

In differential form,

$$d_t f(t,T) = \alpha(t,T)dt + \sigma(t,T) \cdot dW_t.$$

For example, we may have

$$d_t f(t,T) = \alpha(t,T)dt + \sigma_1 dW_t^1 + \sigma_2 e^{-\kappa(T-t)} dW_t^2,$$

where σ_1 , σ_2 , and κ are constants. In this model, W^1 provides "shocks" that are felt equally by all points on the yield curve and W^2 "shocks" that are felt only in the short term.

For a small interval Δ ,

$$f(t+\Delta,T) - f(t,T) \approx \Delta \alpha(t,T) + \sum_{i=1}^{d} \sigma_i(t,T) (W_{t+\Delta}^i - W_t^i).$$

Hence

$$\lim_{\Delta \to 0} \frac{1}{\Delta} \operatorname{var} \left(f(t + \Delta, T) - f(t, T) \right) = \sum_{i=1}^{d} \sigma_i^2(t, T)$$

And

$$\lim_{\Delta \to 0} \frac{1}{\Delta} \operatorname{cov} \left(f(t + \Delta, T) - f(t, T), f(t + \Delta, S) - f(t, S) \right) = \sum_{i=1}^{d} \sigma_i(t, T) \sigma_i(t, S).$$

We may define an instantaneous correlation coefficient for the increments of the forward rate,

$$\frac{\sum_{i=1}^{d} \sigma_i(t,T) \sigma_i(t,S)}{\sqrt{\sum_{i=1}^{d} \sigma_i^2(t,T) \cdot \sum_{i=1}^{d} \sigma_i^2(t,S)}}$$

If d = 1, the increments of the forward rates are perfectly correlated everywhere on the yield curve.

The results of single-factor HJM may be easily generalized. Again we use M_t as numeraire and denote the discounted bond price as $Z(t,T) = M_t^{-1}P(t,T)$. We have

$$d_t Z(t,T) = Z(t,T) \left[\left(\frac{1}{2} (\Sigma \cdot \Sigma)(t,T) - \int_t^T \alpha(t,u) du \right) dt + \Sigma(t,T) \cdot dW_t \right]$$

Then we seek an $\eta = (\eta_t) \in \mathbb{R}^d$ such that

$$\Sigma(t,T) \cdot \eta_t = \frac{1}{2} (\Sigma \cdot \Sigma)(t,T) - \int_t^T \alpha(t,u) du.$$

For the above to have solution, it is necessary that the matrix $\Sigma \equiv (\Sigma^i(t, T_j))$ be full rank for all t and T_j . The we define a $\tilde{\mathbb{P}}$ such that \tilde{W}_t defined below is $\tilde{\mathbb{P}}$ -BM,

$$dW_t = dW_t + \eta_t dt.$$

Under $\tilde{\mathbb{P}}$, $Z_t(t,T)$ is then martingale,

$$d_t Z(t,T) = Z(t,T)\Sigma(t,T) \cdot d\tilde{W}_t.$$

The bond price satisfies

$$d_t P(t,T) = P(t,T) \left(r_t dt + \Sigma(t,T) \cdot d\tilde{W}_t \right)$$

And the forward rate,

$$d_t f(t,T) = -(\sigma \cdot \Sigma)(t,T)dt + \sigma \cdot d\tilde{W}_t.$$

Multi-factor Short Rate Models

Let $\tilde{W} = (\tilde{W}^1, \tilde{W}^2, ..., \tilde{W}^d)$ be a *d*-dimensional Brownian Motion under $\tilde{\mathbb{P}}$. And Let $X = (X^1, X^2, ..., X^N)$ be the *N* factors that determines the short rate r_t . X generally includes a factor that is the short rate itself.

We write

$$r_t = r(X_t), \tag{4.12}$$

where X_t satisfies

$$dX_t = \nu(X_t)dt + \rho(X_t) \cdot d\tilde{W}_t.$$
(4.13)

The term structure P(t,T) can then be represented as

$$P(t,T) = \tilde{\mathbb{E}}_t \left[\exp\left(-\int_t^T r(X_s) ds\right) \right].$$

Of course, any derivative that has a terminal payment of $g(X_T)$ may be priced as

$$\widetilde{\mathbb{E}}_t\left[\exp\left(-\int_t^T r(X_s)ds\right)g(X_T)\right].$$

Feynman-Kac Formulation

It is easy to extend the Feynman-Kac formulation of single-factor short rate model in (4.11) to the multi-factor case. Let F(x,t) = P(t,T), we have

$$F_2(x,t) + \nu(x) \cdot F_1(x,t) + \frac{1}{2} \operatorname{tr} \left[\rho(x)\rho(x)'F_{11}(x,t)\right] - r(x)F(x,t) = 0, \quad (4.14)$$

with

$$F(x,T) = 1.$$
 (4.15)

Obviously, if we change the boundary condition in (4.15) to F(x,T) = g(x), F(x,t) prices any general derivative with terminal payment $g(X_T)$.

4.5 Pricing Interest Rate Products

In this section we briefly review the pricing of some popular interest rate products, given the term structure P(t, T).

Bond with Fixed Coupons

Suppose the coupon rate (uncompounded) is k and the payment is made at a sequence of dates $T_i = T_0 + i\Delta$. The cash flow is shown in the diagram:



This is equivalent to owning a T_n -bond and $k\Delta$ units of T_i -bond for each i = 1, ..., n:

$$\begin{cases} P(T_0, T_n) \\ k\Delta P(T_0, T_i), \ i = 1, ..., n. \end{cases}$$

From

$$k\Delta \sum_{i=1}^{n} P(T_0, T_i) + P(T_0, T_n) = 1,$$

we can determine the appropriate coupon rate,

$$k = \frac{1 - P(T_0, T_n)}{\Delta \sum_{i=1}^{n} P(T_0, T_i)}.$$

Floating-Rate Bond

Now the coupon rate paid at time T_i is the floating rate at previous payment date T_{i-1} , which is defined as

$$L(T_{i-1}) = \frac{1}{\Delta} \left(\frac{1}{P(T_{i-1}, T_i)} - 1 \right).$$

The cash flow is illustrated in the following diagram:

The value of $\Delta L(T_{i-1})$ at T_0 is

$$M_{T_0}\tilde{\mathbb{E}}_{T_0}\left(M_{T_i}^{-1}\Delta L(T_{i-1})\right) = M_{T_0}\tilde{\mathbb{E}}_{T_0}\left[M_{T_i}^{-1}\left(\frac{1}{P(T_{i-1},T_i)}-1\right)\right]$$

$$= M_{T_0}\tilde{\mathbb{E}}_{T_0}\left[P^{-1}(T_{i-1},T_i)\tilde{\mathbb{E}}_{T_{i-1}}\left(M_{T_i}^{-1}\right)-M_{T_i}^{-1}\right]$$

$$= M_{T_0}\tilde{\mathbb{E}}_{T_0}\left[\tilde{\mathbb{E}}_{T_{i-1}}M_{T_{i-1}}^{-1}-M_{T_i}^{-1}\right]$$

$$= M_{T_0}\tilde{\mathbb{E}}_{T_0}\left[M_{T_{i-1}}^{-1}-M_{T_i}^{-1}\right]$$

$$= P(T_0,T_{i-1})-P(T_0,T_i).$$

Note that $P^{-1}(T_{i-1}, T_i)$ is $\mathcal{F}_{T_{i-1}}$ -measurable and $P(T_{i-1}, T_i) = M_{T_{i-1}} \tilde{\mathbb{E}}_{T_{i-1}} M_{T_i}^{-1}$. The contingent claim $\Delta L(T_{i-1})$ can be replicated by buying a T_{i-1} -bond and sell a T_i -bond. At time T_{i-1} , buy $P^{-1}(T_{i-1}, T_i)$ units of T_i -bond. The value of floating rate bond is then

$$P(T_0, T_n) + \sum_{i=1}^n \left(P(T_0, T_{i-1}) - P(T_0, T_i) \right) = P(T_0, T_0) = 1.$$

Swaps

A swap contract exchanges a sequence of floating rate payments for a sequence of fixed-rate payments or vice sersa. The cash flow for the party receiving fixed-rate payments is shown in the following diagram:



Entering a swap contract is equivalent to buying a fixed-coupon bond and selling a floating-rate bond. The former is worth

$$P(T_0, T_n) + k\Delta \sum_{i=1}^{n} P(T_0, T_i),$$

and the latter is worth 1. So the fixed coupon rate k must satisfy

$$k = \frac{1 - P(T_0, T_n)}{\Delta \sum_{i=1}^{n} P(T_0, T_i)}.$$

Forward Swaps

The value of the swap at time T_0 is

$$X = P(T_0, T_n) + k\Delta \sum_{i=1}^{n} P(T_0, T_i) - 1.$$

The value of X at time $t < T_0$ must be

$$M_{t}\tilde{\mathbb{E}}_{t}\left(M_{T_{0}}^{-1}X\right) = M_{t}\tilde{\mathbb{E}}_{t}\left(M_{T_{0}}^{-1}P(T_{0},T_{n}) + k\Delta\sum_{i=1}^{n}M_{T_{0}}^{-1}P(T_{0},T_{i}) - M_{T_{0}}^{-1}\right)$$

$$= M_{t}\tilde{\mathbb{E}}_{t}\left(\tilde{\mathbb{E}}_{T_{0}}M_{T_{n}}^{-1} + k\Delta\sum_{i=1}^{n}\tilde{\mathbb{E}}_{T_{0}}M_{T_{i}}^{-1} - M_{T_{0}}^{-1}\right)$$

$$= M_{t}\tilde{\mathbb{E}}_{t}M_{T_{n}}^{-1} + k\Delta\sum_{i=1}^{n}M_{t}\tilde{\mathbb{E}}_{t}M_{T_{i}}^{-1} - M_{t}\tilde{\mathbb{E}}_{t}M_{T_{0}}^{-1}$$

$$= P(t,T_{n}) + k\Delta\sum_{i=1}^{n}P(t,T_{i}) - P(t,T_{0}) = 0.$$

So the forward swap rate must be

$$k = \frac{P(t, T_0) - P(t, T_n)}{\Delta \sum_{i=1}^{n} P(t, T_i)}.$$

When $t = T_0$, k is equal to the swap rate.

Swaptions

A swaption is a contract to enter a swap at time T_0 with swap rate k. The value of the swaption at time $t < T_0$ is

$$M_t \tilde{\mathbb{E}}_t \left(M_{T_0}^{-1} \max\left(P(T_0, T_n) + k\Delta \sum_{i=1}^n P(T_0, T_i) - 1, 0 \right) \right).$$

The pricing of such swaptions requires the joint distribution of $L(T_i)$ $(0 \le i < n)$ under a single probability measure. We will come back to this problem in the next chapter.

Caps and Floors

A caps contract is an agreement that never pays more than a fixed rate k. So a caps contract pays at time T_i , $1 \le i \le n$,

$$c_i = \Delta \max \left(L(T_{i-1}) - k, 0 \right).$$

For each i, c_i is called a *caplet*. In contrast, a floors contract pays at T_i

$$f_i = \Delta \max \left(k - L(T_{i-1}), 0 \right).$$

For each i, f_i is called a *flootlet*. Obviously, each caplet (floorlet) is a call (put) option on a forward rate.

Note that

$$\begin{aligned} \Delta \left(L(T_{i-1}) - k \right) &= \frac{1}{P(T_{i-1}, T_i)} - 1 - \Delta k \\ &= \left(1 + \Delta k \right) P^{-1}(T_{i-1}, T_i) \left(\frac{1}{1 + \Delta k} - P(T_{i-1}, T_i) \right). \end{aligned}$$

So the value of c_i at time t is

$$M_{t}\tilde{\mathbb{E}}_{t}\left(M_{T_{i}}^{-1}c_{i}\right)$$

$$= (1 + \Delta k)M_{t}\tilde{\mathbb{E}}_{t}\left(M_{T_{i}}^{-1}P^{-1}(T_{i-1}, T_{i})\max\left(\frac{1}{1 + \Delta k} - P(T_{i-1}, T_{i}), 0\right)\right)$$

$$= (1 + \Delta k)M_{t}\tilde{\mathbb{E}}_{t}\left(\left(\tilde{\mathbb{E}}_{T_{i-1}}M_{T_{i}}^{-1}\right)P^{-1}(T_{i-1}, T_{i})\max\left(\frac{1}{1 + \Delta k} - P(T_{i-1}, T_{i}), 0\right)\right)$$

$$= (1 + \Delta k)M_{t}\tilde{\mathbb{E}}_{t}\left(M_{T_{i-1}}^{-1}\max\left(\frac{1}{1 + \Delta k} - P(T_{i-1}, T_{i}), 0\right)\right).$$

The third equality uses the fact that $P(T_{i-1}, T_i) = M_{T_{i-1}} \mathbb{E} M_{T_i}^{-1}$. So a caplet is equivalent to $(1 + \Delta k)$ units of put option on the T_i -bond with strike price $(1 + \Delta k)$ and maturity date T_{i-1} . A floorlet can be similarly interpreted.

4.6 Forward Measure

Recall that the pricing formula,

$$x_t = M_t \tilde{\mathbb{E}}_t M_T^{-1} X_T,$$

where X_T is a contingent claim to be realized at time T and the expectation is taken with respect to the risk-neutral probability measure $\tilde{\mathbb{P}}$. To compute the price, we need the joint distribution of M_T and X_T under $\tilde{\mathbb{P}}$. This is sometimes not convenient for practitioners. In this section, we study forward measure, under which we may obtain simpler pricing formula for many derivatives.

Note that with $M_0 = 1$, we have $P(0,T) = \tilde{\mathbb{E}}M_T^{-1}$. If we define

$$\xi_T = \frac{1}{P(0,T)M_T} = \frac{M_T^{-1}}{\tilde{\mathbb{E}}M_T^{-1}},\tag{4.16}$$

we have $\xi_T > 0$ a.s. and $\mathbb{E}\xi_T = 1$. Hence the \mathcal{F}_T -measurable random variable ξ_T can be the Radon-Nikodym derivative for an equivalent measure for \mathbb{P} . Hence we define an equivalent measure \mathbb{Q}^T as follows,

$$d\mathbb{Q}^T = \xi_T d\tilde{\mathbb{P}}.\tag{4.17}$$

We call \mathbb{Q}^T the *T*-forward measure. Obviously, as $T \to 0$, \mathbb{Q}^T reduces to $\tilde{\mathbb{P}}$. The density process associated with ξ_T is given by

$$\xi_t = \tilde{\mathbb{E}}_t \xi_T = \frac{1}{P(0, T)M_t} \left(M_t \tilde{\mathbb{E}}_t M_T^{-1} \right) = \frac{P(t, T)}{P(0, T)M_t}.$$
(4.18)

Indeed, as shown in Section 4.2, ξ_t is an exponential martingale with $\xi_0 = 1$ and

$$d\xi_t = \xi_t \Sigma(t, T) d\tilde{W}_t.$$

We have

$$\xi_t = \exp\left(\int_0^t \Sigma(s,T) d\tilde{W}_s - \frac{1}{2} \int_0^t \Sigma^2(s,T) ds\right).$$

Define a process W_t^T that satisfies $W_0^T = 0$ a.s. and

$$dW_t^T = d\tilde{W}_t - \Sigma(t, T)dt.$$
(4.19)

By Girsanov Theorem, W_t^T is a BM under \mathbb{Q}^T .

Pricing with Forward Measure Using the forward-T measure, the no-arbitrage price of the contingent claim X_T is given by

$$\begin{aligned} x_t &= M_t \mathbb{E}_t M_T^{-1} X_T \\ &= P(t,T) \xi_t^{-1} \tilde{\mathbb{E}}_t \xi_T X_T \\ &= P(t,T) \mathbb{E}_t^T X_T, \end{aligned}$$
(4.20)

where \mathbb{E}_t^T is taken under \mathbb{Q}^T . Note that P(t,T) is observable at t and that we only need the distribution of X_T under \mathbb{Q}^T to calculate the current price of X_T .

In general, the no-arbitrage price of a contingent claim X_S with $S \in (t, T]$ is

$$x_{t} = M_{t} \tilde{\mathbb{E}}_{t} M_{S}^{-1} X_{S}$$

$$= P(t,T) \xi_{t}^{-1} \tilde{\mathbb{E}}_{t} \xi_{S} \frac{X_{S}}{P(S,T)}$$

$$= P(t,T) \mathbb{E}_{t}^{T} \frac{X_{S}}{P(S,T)}.$$
 (4.21)

This confirms that, to preclude arbitrage opportunity, asset prices with P(t,T) as the numeraire must be martingales under the forward-T measure.

The Forward Rate Under \mathbb{Q}^T , the expectation hypothesis holds:

$$f(t,T) = -\frac{\partial}{\partial T} \log P(t,T)$$

$$= -\frac{1}{P(t,T)} \frac{\partial}{\partial T} \tilde{\mathbb{E}}_t \exp\left(-\int_t^T r_s ds\right)$$

$$= \frac{M_t}{P(t,T)} \tilde{\mathbb{E}}_t M_T^{-1} r_T$$

$$= \xi_t^{-1} \tilde{\mathbb{E}}_t \xi_T r_T$$

$$= \mathbb{E}_t^T r_T,$$

where ξ_t is given in (4.18). Note that, in general, the expectation hypothesis does not hold under \mathbb{P} .

We may further infer that f(t,T) is a martingale under \mathbb{Q}^T . To see this, note that $r_T = f(T,T)$ and that, using (4.19), we have

$$d_t f(t,T) = -\sigma(t,T)\Sigma(t,T)dt + \sigma(t,T)d\tilde{W}_t$$

= $\sigma(t,T)dW_t^T$,

where $\Sigma(t,T) = -\int_t^T \sigma(t,u) du$.

Other Forward Measures With P(t,T) as numeraire, prices of discount bonds with other maturities, say P(t,S) with S < T, are martingales under \mathbb{Q}^T . That is, P(t,S)/P(t,T) is a \mathbb{Q}^T -martingale for any 0 < S < T. In fact, we can prove that

$$\frac{P(t,S)}{P(t,T)} = \frac{P(0,S)}{P(0,T)} \exp\left(L_t - \frac{1}{2}[L]_t\right),$$
(4.22)

where $L_t = -\int_0^t \Sigma_{S,T}(s) dW_s^T$ with $\Sigma_{S,T}(t) \equiv \Sigma(t,T) - \Sigma(t,S) = -\int_S^T \sigma(t,u) du$. To show this, denote $Y(t,S) \equiv P(t,S)/M_t$ and $Z(t,T) \equiv P(t,T)/M_t$. We have P(t,S)/P(t,T) = Y(t,S)/Z(t,T) and, from (4.3),

$$\begin{aligned} &d_t Y(t,S) = Y(t,S)\Sigma(t,S)d\tilde{W}_t \\ &d_t Z(t,T) = Z(t,T)\Sigma(t,T)d\tilde{W}_t. \end{aligned}$$

Applying Ito's lemma, we have

$$\begin{aligned} d_t \frac{Y(t,S)}{Z(t,T)} &= \frac{1}{Z(t,T)} d_t Y(t,S) - \frac{Y(t,S)}{Z^2(t,T)} d_t Z(t,T) + \frac{Y(t,S)}{Z^3(t,T)} d[Z]_t - \frac{1}{Z^2(t,T)} d[Z,Y]_t \\ &= -\frac{Y(t,S)}{Z(t,T)} \Sigma_{S,T}(t) \left(d\tilde{W}_t - \Sigma(t,T) dt \right) \\ &= -\frac{Y(t,S)}{Z(t,T)} \Sigma_{S,T}(t) dW_t^T. \end{aligned}$$

We thus obtain

$$d_t \frac{P(t,S)}{P(t,T)} = \frac{P(t,S)}{P(t,T)} dL_t,$$

which is the differential form of (4.22).

Using (4.22), we can define S-forward measure \mathbb{Q}^S with the density process

$$\xi_t = \frac{P(0,T)}{P(0,S)} \frac{P(t,S)}{P(t,T)} = \exp\left(L_t - \frac{1}{2}[L]_t\right).$$
(4.23)

Hence we obtain a spectrum of equivalent martingale measures, each of which corresponds to the numeraire of P(0, S).

Pricing Bond Options Consider a risk-free discount bond maturing on T and a European call option on the bond with expiration date $S \leq T$ and strike price K. Using (4.21), the price of the call at time 0 is given by

$$P(0,T)\mathbb{E}^{T} \left[P(S,T)^{-1} \max(P(S,T) - K,0) \right]$$

= $P(0,T)\mathbb{E}^{T} I\{P(S,T) \ge K\} - KP(0,T)\mathbb{E}^{T} \left[P(S,T)^{-1} I\{P(S,T) \ge K\} \right]$
= $P(0,T)\mathbb{Q}^{T} \{P(S,T) \ge K\} - KP(0,S)\xi_{0}^{-1}\mathbb{E}^{T} \left[\xi_{S} I\{P(S,T) \ge K\}\right]$
= $P(0,T)\mathbb{Q}^{T} \{P(S,T) \ge K\} - KP(0,S)\mathbb{Q}^{S} \{P(S,T) \ge K\},$

where ξ_t is defined in (4.23). Note that

$$\mathbb{Q}^{T} \{ P(S,T) \ge K \} = \mathbb{Q}^{T} \left\{ \frac{P(S,S)}{P(S,T)} \le \frac{1}{K} \right\} = \mathbb{Q}^{T} \left\{ \log \frac{P(S,S)}{P(S,T)} \le -\log K \right\}$$
$$\mathbb{Q}^{S} \{ P(S,T) \ge K \} = \mathbb{Q}^{S} \left\{ \frac{P(S,T)}{P(S,S)} \ge K \right\} = \mathbb{Q}^{S} \left\{ \log \frac{P(S,T)}{P(S,S)} \ge \log K \right\}.$$

Note that $\frac{P(t,S)}{P(t,T)}$ and $\frac{P(t,T)}{P(t,S)}$ are martingales under \mathbb{Q}^T and \mathbb{Q}^S , respectively. We can show (left as an exercise) that

$$\log \frac{P(S,S)}{P(S,T)} = \log \frac{P(0,S)}{P(0,T)} - \frac{1}{2} \int_0^S \Sigma_{S,T}(u)^2 du - \int_0^S \Sigma_{S,T}(u) dW_u^T,$$

so that

$$\log \frac{P(S,S)}{P(S,T)} \sim N\left(\log \frac{P(0,S)}{P(0,T)} - \frac{1}{2} \int_0^S \Sigma_{S,T}^2(u) du, \int_0^S \Sigma_{S,T}^2(u) du\right).$$

So $\frac{P(S,S)}{P(S,T)}$ is log normal if $\sigma(t,T)$ (and thus $\Sigma(t,T)$) is deterministic. The Vasicek model, for example, has $\sigma(t,T) = \sigma \exp(\alpha(T-t))$. Similarly, we can obtain the distribution of $\log \frac{P(S,T)}{P(S,S)}$ under \mathbb{Q}^S . It turns out that the pricing formula of the European call on the T-bond is given by

$$P(0,T)\Phi(d_1) - KP(0,S)\Phi(d_2),$$

where Φ is the cumulative distribution function of N(0, 1) and

$$d_{1,2} = \frac{\log \frac{P(0,T)}{KP(0,S)} \pm \frac{1}{2} \int_0^S \Sigma_{S,T}^2(u) du}{\left(\int_0^S \Sigma_{S,T}^2(u) du\right)^{1/2}}.$$

Appendix A

Appendix to Chapter 1

A.1 Classical Derivation of CAPM

A.1.1 Efficiency Frontier without Riskfree Asset

Suppose there are N risky assets, and the return vector of these assets has a mean of μ and a covariance matrix Σ . Each agent selects a portfolio of these assets h, where $h'\iota = 1$ and ι is a vector of ones. Thus the portfolio return has mean $\mu_h = h'\mu$ and variance $\sigma_h^2 = h'\Sigma h$. We assume that all agents in the economy are identical with utility function $u(\mu_h, \sigma_h^2)$. It is understood that $u(\cdot, \cdot)$ is increasing in μ_h and decreasing in σ_h^2 .

Given an objective mean return of portfolio, agents try to find a portfolio that minimizes the variance. Mathematically, the following problem is to be solved,

$$\min_{h} \frac{1}{2}h'\Sigma h,$$

subject to

$$h'\mu = \mu_h$$
 and $h'\iota = 1$.

The Lagrangian function is given by

$$L = \frac{1}{2}h'\Sigma h + \lambda_1(h'\mu - \mu_h) + \lambda_2(h'\iota - 1).$$

The first order conditions are

$$\Sigma h = \lambda_1 \mu + \lambda_2 \iota, \tag{A.1}$$

$$\mu'h = \mu_h, \tag{A.2}$$

$$\iota'h = 1. \tag{A.3}$$

(A.1) yields

$$h = \Sigma^{-1} (\lambda_1 \mu + \lambda_2 \iota). \tag{A.4}$$

Pre-multiply (A.4) with μ' and ι' , respectively, we obtain

$$\begin{aligned} A\lambda_1 + B\lambda_2 &= \mu_h \\ B\lambda_1 + C\lambda_2 &= 1, \end{aligned}$$

where

$$A = \mu' \Sigma^{-1} \mu, \quad B = \mu' \Sigma^{-1} \iota, \quad C = \iota' \Sigma^{-1} \iota.$$

Define $D = AC - B^2$. We obtain

$$\lambda_1 = \frac{1}{D}(C\mu_h - B), \quad \lambda_2 = \frac{1}{D}(-B\mu_h + A).$$

Plug in (A.4), we obtain

$$h = g_0 + g_1 \mu_h, \tag{A.5}$$

where

$$g_0 = \frac{1}{D} (A \Sigma^{-1} \iota - B \Sigma^{-1} \mu), \quad g_1 = \frac{1}{D} (C \Sigma^{-1} \mu - B \Sigma^{-1} \iota).$$

The variance of the return on h is given by

$$\sigma_h^2 = h' \Sigma h = g_0' \Sigma g_0 + 2\mu_h g_0' \Sigma g_1 + \mu_h^2 g_1' \Sigma g_1.$$

Hence the pairs (μ_h, σ_h) trace a hyperbola boundary, the upper boundary of which is called the *efficiency frontier*.

Note that the minimum-variance portfolio h (A.5) is linear in μ_h . If we know two minimum-variance portfolios h_1 and h_2 with mean returns μ_1 and μ_2 , respectively, then we know all minimum-variance portfolios. Indeed, for all expected return μ_a , the corresponding minimum-variance portfolio can be constructed by $h_a = \alpha h_1 + (1 - \alpha)h_2$, where α is obtained by solving $\mu_a = \alpha \mu_1 + (1 - \alpha)\mu_2$. This observation is often called the *two mutual fund theorem*.

A.1.2 CAPM

Suppose there is a money account with a risk-free return of r. Now the agents' problem becomes

$$\min_{h} \frac{1}{2}h' \Sigma h,$$

subject to

$$\mu' h + (1 - \iota' h)r = \mu_h.$$

The Lagrangian is

$$L = \frac{1}{2}h'\Sigma h + \lambda(\mu_h - \mu'h - (1 - \iota'h)r).$$

The first-order conditions are

$$\Sigma h = \lambda(\mu - r\iota)$$
$$\mu' h + (1 - \iota' h)r = \mu_h.$$

Solving this set of equations, we obtain

$$h = \frac{\mu_h - r}{(\mu - r\iota)' \Sigma^{-1} (\mu - r\iota)} \cdot \Sigma^{-1} (\mu - r\iota).$$
 (A.6)

Note that this portfolio equals a scalar that depends on μ_h times a vector that does *not* depend on μ_h . In other words, for all expected return, the exactly same proportion of each risky assets are chosen. We normalize $\Sigma^{-1}(\mu - r\iota)$ to obtain the so-called tangency portfolio,

$$h_m = \frac{\Sigma^{-1}(\mu - r\iota)}{\iota'\Sigma^{-1}(\mu - r\iota)}.$$

The normalization makes the elements in h_m adding up to one. In the idealized world of CAPM, therefore, everyone will choose the tangency portfolio. Individuals differ only in the percentage of cash holding or leverage. In equilibrium, the *market portfolio* must be the tangency portfolio.

Let R_m be the market return. The variance of the market return is given by

$$\operatorname{var}(R_m) = h'_m \Sigma h_m = \frac{(\mu - r\iota)' \Sigma^{-1} (\mu - r\iota)}{(\iota' \Sigma^{-1} (\mu - r\iota))^2}.$$

The expected market premium over the risk-free return,

$$\mathbb{E}R_m - r = h'_m \mu - r = \frac{(\mu - r\iota)'\Sigma^{-1}(\mu - r\iota)}{\iota'\Sigma^{-1}(\mu - r\iota)}.$$

Let R_i be the *i*-th asset. We have $\mathbb{E}R_i = e'_i\mu$ and

$$\operatorname{cov}(R_i, R_m) = e_i' \Sigma h_m$$

where e_i is a vector that has 1 on the *i*-th element and 0 on others. Now we have

$$(\mathbb{E}R_m - r)\frac{\operatorname{cov}(R_i, R_m)}{\operatorname{var}(R_m)} = e'_i(\mu - r\iota) = \mathbb{E}R_i - r.$$

Rearranging terms, we obtain the celebrated CAPM model,

$$\mathbb{E}R_i = r + \beta_i (\mathbb{E}R_m - r), \tag{A.7}$$

where

$$\beta_i = \frac{\operatorname{cov}(R_i, R_m)}{\operatorname{var}(R_m)}.$$
(A.8)

In the CAPM model, the expected payoff of a security is a linear function of the security's beta, which characterizes the systematic risk contained in the security. The linear function is called the *security market line* (SML). Obviously, the intercept of the SML is the risk-free rate (r) and the slope of the SML is the market risk premium $(\mathbb{E}R_m - r)$.

Index

adapted, 14 affine model, 76 arbitrage, 43 Bayes rule, 51 Black-Karasinski model, 75 Black-Scholes model, 44 bond float rate, 80 with fixed coupons, 80 Brownian motion, 15 Brownian motion with drift, 26 caps, 83 cash flow pricing, 56 change of measure, 49 constant-elasticity diffusion, 26 Cox-Ingersoll-Ross model, 75 delta, 44 delta hedging, 45 density process, 51 diffusion, 24 diffusion function, 24 drift function, 24 equivalent probability measure, 50 Euler approximation, 29 event, 13 exponential martingale, 24 Feller's squared-root Process, 27 Feynman-Kac solution, 46 Feynman-Kac solution multivariate case, 48

filtration, 14 floors, 83 forward price of bond, 67 forward rate, 67 forwards, 54 forwards-futures spread, 55 futures, 55 gains process, 42 gamma, 44 geometric Brownian motion, 26 geometric form, 42 Girsanov multivariate, 56 Girsanov theorem, 52 Girsanov theorem light version, 52 Heath-Jarrow-Morton model multi-factor, 77 single factor, 68 Ho and Lee model, 72 integration by parts, 22 Ito integral, 19 Ito's formula, 22, 23 Ito's formula multivariate, 23 linear drift, 25 market price of risk, 54 Markov process, 16 Markov process homogenous, 17

martingale, 15 martingale equivalent measure, 50 Milstein approximation, 30 natural filtration, 14 numeraire, 43 numeraire invariance theorem, 43 Ornstein-Uhlenbeck process, 27 portfolio, 42 quadratic covariation, 20 quadratic variation, 20 random variable, 13 self-financing, 43 semimartingale, 20 short rate, 66 short rate model, 71 sigma field, 13 sigma field generated by a random variable, 14 spot rate, 65 Stieltjes, 19 stochastic process, 14 Stratonovich integral, 19 sub-martingale, 15 sup-martingale, 15 swaps, 81 swaptions, 82 term structure, 65 term to maturity (TTM), 66 theta, 44 trading strategy, 42 transition density, 17 transition probability, 16 Vasicek model, 74 yield curve, 66 yield to maturity (YTM), 66